arXiv:1409.5953v1 [math.GR] 21 Sep 2014

ITERATED IDENTITIES AND ITERATIONAL DEPTH OF GROUPS

ANNA ERSCHLER

ABSTRACT. Given word on n letters, we study groups which satisfiy "iterated identity" w, meaning that for all $x_1, \dots, x_n$ there exists m such that m-the iteration of w of Engel type, applied to $x_1, \dots, x_n$, is equal to the identity. We define bounded groups and groups which are fractal with respect to identities. This notion of being fractal can be viewed as a self-similarity conditions for the set of identities, satisfied by a group. In contrast with torsion groups and Engel groups, groups which are fractal with respect to identities appear among finitely generated elementary amenable groups. We prove that any polycyclic, as well as any metabelian group is bounded and we compute the iterational depth for various wreath products. We study the set of iterated identities, satisfied by a given group, which is not necessarily a subgroup of a free group and not necessarily invariant under conjugation, in contrast with usual identities. Finally, we discuss another notion of iterated identities of groups, which we call solvability type iterated identities, and its relation to elementary classes of varieties of groups.

1. INTRODUCTION

Given a word $w = w(x_1, \dots, x_n)$ on n letters, we recall that a group G satisfies an identity w if for any $y_1, \dots, y_n \in G$ it holds $w(y_1, \dots, y_n)$. A well-known identity on one single letter is $w(x) = x^p$. For p being large enough, it is known that there are infinite finitely generated groups satisfying this identity: that is, infinite finitely generated groups such that $g^p = e$ for any $g \in G$. A larger class of groups consists of groups that are p-groups, that is, G are such that for any $g \in G$ there exists k such that $g^{p^k} = e$. A first group of this kind was constructed by Golod ([12]), further examples include Grigorchuk group [13], Gutpa-Sidki groups [17]); a large family of p-torsion groups of unbounded torsion appear as quotient groups of hyperbolic groups (see Gromov [16]). The groups above do not satisfy the identity $w = x^p$, that is, there exist $g \in G$ such that $w(g) \neq e$, but they satisfy some iteration of w: for any $g \in G$ there exists k such that $w(w(\dots w(g) \dots)) = e$.

What happens with other identities w if we consider groups that satisfy some iteration of w? $w(x) = x^k$ being the only identity in one letter, so we are mostly interested in the case where $n \geqslant 2$. In this situation, in contrast with one letter case, there are several possible notion of "iterated identity", some of which we discuss in this paper.

Before giving the definitions, we mention possible motivation to study iterated identities of groups:

In the case when a group satisfying an identity w are well-understood, the groups satisfying an iterated identity w can be much more complicated: consider e.g. $w(x_1, x_2) = x_1 x_2 x_1^{-1} x_2^{-1}$; groups satisfying w are Abelian, whereas the structure of Engel groups is far from being understood; in the case when groups satisfying an identity are already tricky, like Burnside groups satisfying an identity $w = x^p$, the groups satisfying the iteration of the identity form a larger, and possibly even trickier, class of groups.

The work is supported by the ERC grant 257110 "RaWG" and the ANR "DiscGroup: facettes des groupes discrets" .

Moreover, it might be possible to construct groups satisfying additional natural properties among groups satisfying iterated identities, such that groups satisfying the same identity do not allow: for example, there exist *residually finite* infinite finitely generated torsion groups ([4, 12, 13, 17],) and residually finite infinite finitely generated non-nilpotent Engel group (Golod, [12]), while any finitely generated residually finite torsion group of bounded torsion is finite (follows the solution by Zelmanov of so called restricted Burnside problem [30]); any finitely generated residually finite group that satisfies a fixed bounded iteration of the identity $ab = ba$ is nilpotent (Wilson, [29], Theorem 2, the proof uses classification of finite simple groups).

Given a group G, it seems interesting to describe all iterated identities that G satisfies. It may happen that the group does not satisfy any non-trivial identity, but does satisfy non-trivial iterated identities (e.g. first Grigorchuk group and many other groups acting on a rooted tree, does not satisfy any identity by a result of Abert [1]; yet it is well known that this group is a torsion 2 group, that is, satisfy the iteration of $w(x_1) = x_1^2$ [13]). The set of iterated identities on n of variables does not necessarily form a subgroup of the free group F_n (see Example 2.7) and it is not necessarily invariant with respect to conjugation (see Example 4.4), in contrast with usual identities, which is one of the difficulties to study this set. There are various invariants of groups which can be defined in terms of the set of all iterated identities satisfied by a group, such as boundedness and fractalness with respect to identities, which we discuss in this paper.

In some situations iterated identities and iterated quasi-varieties provide rougher invariants, for example there are various iterated identities, satisfied by groups which do not satisfy any non-trivial identity. In other situations iterated identities and iterated quasi-varieties can provide softer invariants: it may happen that two groups G and H that generate distinct varieties of groups, but iterated quasi-varities generated by these groups (both Engel type iterated quasi-varieties for a fixed (small) number of variables which hold in a given group as well as solvability type iterated quasi-varities, which we define in Section 8) are the same;

Let $w = w(x_1, \dots, x_n)$ be a word on n letters, $n \geqslant 1$.

Definition 1.1. We say that a group G satisfies E-type (or Engel type) iterated identity w if for any $x_1, \dots, x_n \in G$ there exists N such that n-th iteration of w on the first variable

$$w_{\circ N}(x_1, \dots, x_n) = w(w(\dots(w(x_1, x_2, \dots, x_n), x_2, \dots, x_n), x_2, \dots, x_n)) = e.$$

Some iterated E type identities on two variables are a particular case of a closely related definition of "correct sequences" in terminology of [5, 18, 25]: a sequence $u = u_1, u_2, u_3, \dots, u_n, \dots$ of elements from a free groups on two variables F_2 is said to be *correct* if the following conditions hold: (i) For every group G and elements $a, g \in G$, we have $u_n(a, 1) = e$ and $u_n(1, g) = e$ for all sufficiently large n. (ii) if a, g are elements of G such that $u_n(a, g) = 1$, then for every $m \geqslant n$ it holds $u_m(a, g) = e$.

The words we consier in our definition 1.1 are not necessarily products of commutators, and thus do not need in general to satisfy (ii). It is also essential that in definition 1.1 we do not specify first values of the sequences of iterations.

In [5, 9, 10, 26] one describes some sequences of iteration of an identity which characterize finite solvable groups (we recall these results in subsection 2.1). The situation with infinite groups is essentially different. It is clear for example that no finite set of iterated identities can characterize infinite solvable groups. Instead of searching some iterated identity, we are interested in properties of all iterated identities which hold in a given group or a

class of groups, finite or infinite, and we are mostly interested in finitely generated infinite groups.

It seems particular interesting to descibe iterated identities for some class of groups, that have some other manifestations of self-similarity, such as Grigorchuk groups; Golod groups [12]; Basilica group and other iterated monodromy groups [22]; branch groups [14].

But it seems also interesting to understand the structure of iterated identities for groups without natural self-similarity structure, for example free and not free groups of various varieties.

The questions we want to address: which iterated identities hold for a given group or class of group? Are "quasi-varitety" defined by a set of iterated varieties of two given groups coincide? For which groups it is essential to consider iterated identities rather than identities, that is, in which cases it is essential to allow unbounded number of iteration for a given identity: we say that the iterated identity is bounded if this guaranteed by a bounded number of iteration. We say that the group is bounded, if it has at least one non-trivial iterated identity, all of its iterated identities are bounded and, moreover, the number of iteration for each of them is bounded by a common upper bound (see definitions 3.1 and 3.2 for more details).

The paper has the following structure. Section 2 contains preliminaries about iterated identities. We give first examples of iterated identities. In Example 2.7 we show that a product of iterated identities (in a symmetric group) is not necessarily an iterated identity.

In Section 3 we define iterational depth of an iterated identity and iterational depth of a group, as well as bounded groups and groups which are fractal with respect to identities.

In Section 4 we provide examples of finitely generated elementary amenable groups which are fractal with respect to identities (see Example 4.1). We show that a conjugate of an iterated identity of a finitely generated elementary amenable group is not necessarily an iterated identity of this group (see Example 4.4).

In Section 5 we discuss the behavior of iterational depth with respect to quotients. We show in particular that any polycyclic group is bounded.

In Section 6 we compute the iterational depth of wreath products of cyclic groups. We show that

– The iterational depth of $\mathbb{Z} \wr \mathbb{Z}$ is equal to 2 (see Proposition 6.1)

– The iterational depth of $\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}$ is equal to $k+1$, k being the maximum of α_i, where $R = \prod p_i^{\alpha_i}$, p_i are prime numbers, $\alpha_i \in \mathbb{Z}_+$ (see Proposition 6.3).

We show that a subgroup of a finitely generated bounded group is not always bounded.

In Section 7 we prove

Theorem A (Theorem 7.1)**.** *Any finitely generated metabelian group is bounded.*

In Section 8 we define another notion of iterated identities, which we call solvability type (or S type) iterated identities and we discuss the relation between iterated quasi-varieties and elementary classes generated by varieties of groups. In Example 8.7 we consider first Grigorchuk group and show that a group satisfying an iterated identity $w(x_1) = x_1^2$ does not necessarily belong to the elementary class of groups satisfying bounded iteration of this identity.

In Subsection 8.2 we list some open problems.

1.1. **Aknowledgements.** . The author is grateful to Rostislav Grigorchuk for his remarks on the preliminary version of this paper.

sectionbasic

2. BASIC FACTS AND EXAMPLES

Remark 2.1. Suppose that $w(x_1,\dots,x_n)$ is freely equivalent to $w'(x_1,\dots,x_n)$. Then a group G satisfies an E type iterated identity w if and only if it satisfies an iterated identity w'.

Proof. It is clear that $w_{\circ d}(x_1,x_2,\dots,x_n) = w'_{\circ d}(x_1,x_2,\dots,x_n)$ for any G, any $x_1,\dots,x_n \in G$ and any $d \geqslant 1$. □

Given a group G, we can therefore speak about subset $\Omega_n \subset F_n$ of E type iterated identities of G on n variables. We call this subset Ω_n *characteristic subset* of G.

The following is straightforward

Lemma 2.2. (1) *If G satisfies an E-type iterated identity w, then any subgroup of G has the same property.*
(2) *If G satisfies an E-type iterated identity w, then any quotient of G has the same property.*
(3) *If $G_1 \subset G_2 \subset G_3 \dots$, and each G_i satisfies an E-type identity w, then the union $G = \cup_{i=1}^{\infty} G_i$ satisfies the same property.*

Example 2.3. *Let N be a nilpotent group, $G = N \wr \mathbb{Z}/p\mathbb{Z}$, $w(x_1,x_2) = (x_1x_2x_1^{-1}x_2^{-1})^p$. Then G satisfies an Engel type identity w.*

Proof. First observe that w is an iterated identity of N. Indeed, k-th iteration of w belongs to the k-th group in a central series of N. Observe also that for any $x_2 \in G$ and any $x_1 \in \mathbb{Z}/p\mathbb{Z}^{\infty} \subset N \wr \mathbb{Z}/p\mathbb{Z}$, it holds $(x_1x_2x_1^{-1}x_2^{-1}) \in (\mathbb{Z}/p\mathbb{Z})^{\infty}$, and thus $w(x_1,x_2) = e$. □

examplefreesubgroup

Example 2.4. *i) Let the word $w(x_1,x_2,\dots,x_n)$ be not freely equivalent to the empty word. Then for any $k \geqslant 1$ its iteration $w_{\circ k}(x_1,x_2,\dots,x_n)$ is not freely equivalent to the empty word.*

ii) Let G be a group that contains a non-Abelian free subgroup. Then G does not satisfy any non-trivial E type iterated identity: for any $w(x_1,x_2,\dots,x_n)$ which is not freely equivalent to the empty word, G does not satisfy w.

Proof. To prove i) and ii), it is sufficient to show that the free group F_2 does not satisfy the iterated identity w, if $w(x_1,x_2,\dots,x_n)$ is not freely equivalent to the identity word. First observe that w is not freely equivalent to some word in $x_2,\dots,x_n$.

Indeed, if G is any group satisfying E type iterated identity $w(x_1,x_2,\dots,x_n)$ such that $w(x_1,x_2,\dots,x_n) = w'(x_2,\dots,x_n)$, then all iteration of w are equal to $w'(x_2,\dots,x_n)$. Hence, If w is an E type iterated identity of G, then $w(x_1,x_2,\dots,x_n) = w'(x_2,\dots,x_n)$ is identity of G. If G is a free group, this implies that w is freely equivalent to trivial word.

Now suppose that w is not freely equivalent to some word in $x_2,\dots,x_n$. Since F_2 contains F_n as a subgroup, it suffices to show that w is not an iterated identity of F_n. Let $x_1,\dots,x_n$ be free generators of F_n. Observe that $w(x_1,\dots,x_n)$ does not belong to the subgroup generated by $x_2,\dots,x_n$. Therefore, $w(x_1,\dots,x_n)$, $x_2,\dots,x_n$ generate a free group of rank n. Applying the same argument again, we conclude that any iteration of $w(x_1,\dots,x_n)$ together with $x_2,\dots,x_n$ generate a free group of rank n. In particular, no such iteration is trivial, and hence w is not an iterated identity of G.

□

subsolvable

2.1. **Known results about finite solvable groups.** Below we use the following notation: $w^y = y^{-1}wy$, $w^{-y} = y^{-1}w^{-1}y$, $[u,w] = u^{-1}u^w = u^{-1}w^{-1}uw$, $[w_1,w_2,\dots w_k] = [w_1,w_2],w_3\dots]$.

One of the examples constructed by Brandl and Wilson show in [9] is a sequence of words in 4 letters X, Y, Z, T: $s_1(X,Y,Z,T) = Y$, s_{k+1} is defined recursively by $s_{k+1}(X,Y,Z,T) = [X, s_k(X,Y,Z,T), s_k(X,Y,Z,T), Z]^T$. A sequence of words s_n in variables x, y is defined by Bray, Wilson and Wilson in [10] by the rules $s_0 = x$ and $s_{n+1} = \left[s_n^{-y}, s_n\right]$. Bandman,Greuel, Grunewald, Kunyavskiĭ, Pfister, and Plotkin study in ([5]) the sequence $s_1 = x^{-2}y^{-1}x$ and $s_{n+1} = [xs_n^{-1}x^{-1}, ys_n^{-1}y^{-1}]$ (in [5] the notation used for the commutator $[u,w] = uwu^{-1}w^{-1}$, different from the one we use and from that of the [9, 10, 26]). Ribnere constructs in [26] various sequences of the form $s_{k+1} = [s_k^f, s_k^g]$.

In all these examples it is clear that for any solvable group of solvability length at most k we have $s_k = e$. The results of above mentioned papers is that for finite groups the converse statement holds: any finite group such that $s_k = e$ for any choice of variables and any sufficiently large k is solvable. (The idea of the proofs is that if there would exist a non-solvable finite group G with this property, then we could chose G with this property among minimal simple finite groups. The authors use then the characterization of minimal finite groups due to Thompson ([27, 28] to find a simple group which violate this property and to get a contradiction).

remarkgolod **Remark 2.5.** In all the examples above, it is clear that there exist (infinite) finitely generated non-solvable groups G such that $s_k = e$ for all sufficiently large k. Indeed, if G is such that any 4 generated group of G is solvable (e.g. G is one of Golod groups constructed in [12]), then it is clear that G has this property.

reformulationsolvable **Remark 2.6.** Put

$$w_{\mathrm{BW}}(x_1,x_2,x_3,x_4) = [x_2,x_1,x_1,x_3]^{x_4};$$
$$w_{\mathrm{BWW}}(x_1,x_2) = [x_1^{-x_2},x_1] = [x_2^{-1}x_1^{-1}x_2,x_1] = x_2^{-1}x_1x_2x_1^{-1}x_2^{-1}x_1^{-1}x_2x_1;$$
$$w_{\mathrm{BGGKPP}}(x_1,x_2,x_3) = [x_2x_1^{-1}x_2^{-1}, x_3x_1^{-1}x_3^{-1}] = x_2x_1x_2^{-1}x_3x_1x_3^{-1}x_2x_1^{-1}x_2^{-1}x_3x_1^{-1}x_3^{-1}$$

The results above imply in our terminology that a finite group satisfies an E type iterated identity w_{BW}, (or w_{BWW}, or w_{BGGKPP} etc) if and only if it is solvable; and it is clear that any solvable group, finite or infinite, satisfy these identities. Observe however that a class of (infinite) groups satisfying the iterated identity w_{BGGKPP} might be smaller than the class of groups, satisfying the corresponding correct sequence from [5], cited above.

2.2. **Symmetric groups.**

The following example shows that a product of E type iterated identities is not necessarily an E type iterated identity.

examplenotproduct **Example 2.7.** *Consider a finite symmetric group S_n. Let m be the product of prime numbers that are not greater than n. Suppose that $n \geqslant 6$. Put $w_1(x_1,x_2,x_3,x_4) = x_2x_1^mx_2^{-1}$, $w_2(x_1,x_2,x_3,x_4) = x_3x_1^mx_3^{-1}$, $w_3(x_1,x_2,x_3,x_4) = x_4x_1^mx_4^{-1}$. Then w_1, w_2, w_3 are iterated identities of S_n, while their product $w = w_1w_2w_3$ is not an iterated identity of S_n.*

Proof. Observe that for any $g \in S_n$ there exists $k \geqslant 1$ such that $g^{m^k} = e$. Observe that $w_{1,\circ d}(x_1,x_2,x_3,x_4) = x_2^dx_1^{m^d}x_2^{-d}$, and hence w_1 (and, analogously, w_2 and w_3) is an iterated identity of S_n.

Now we prove that w is not at iterated identity of S_n. The elements of the set of cardinality n we number by 1, 2, …, n. Consider $x_1 \in S_n$ which is a product of a cycle of length four: $1 \to 2 \to 3 \to 4 \to 1$ and of length two: $5 \leftrightarrow 6$. Note that m is an even number which is not divided by four, and hence $g = x_1^m$ is a product of two cycles of length two: $1 \to 3 \to 1$ and $2 \to 4 \to 1$. Observe that x_1 is an even permutation. Let us show that there exist $x_2, x_3, x_4 \in S_n$ such that $x_1 = x_2gx_2^{-1}x_3gx_3^-1x_4gx_4^{-1}$. Indeed, x_1

is a product of the following cycles of length two: $5 \leftrightarrow 6$, $1 \leftrightarrow 2$, $1 \leftrightarrow 3$, $1 \leftrightarrow 4$ (here and in the sequel we multiply the cycles from left to right). Therefore, It is also a product of the following cycles of length two

$$x_1 = ((5 \leftrightarrow 6), (1 \leftrightarrow 2))\,((1 \leftrightarrow 3), (5 \leftrightarrow 6))\,((5 \leftrightarrow 6)(1 \leftrightarrow 4))\,.$$

Observe that a product of two cycles on any two pairs of four disjoint elements is conjugate to g in S_n. We have therefore shown that there exists $x_2, x_3, x_4 \in S_n$ such that

$$x_1 = x_2 x_1^m x_2^{-1} x_3 x_1^m x_3^{-1} x_2 x_1^m x_2^{-1}.$$

In other words, $x_1 = w(x_1, x_2, x_3, x_4)$, and hence $w_{\circ d}(x_1, x_2, x_3, x_4) = x_1 \neq e$ for any $d \geqslant 1$. This shows that w is not an iterated identity of S_n.

□

3. Bounded groups, fractal groups, iterational depth of a group

sectionbounded

defidbounded

Definition 3.1. Given an Engel type iterated identity w of a group G let us say that G has *iterational depth* at most D with respect to w if for any $x_1, \dots, x_n \in G$ there exists $d \leqslant D$ such that d-th Engel type iteration $w_{\circ d}(x_1, \dots, x_n) = e$ in G. The infimum of such D we call *iterational depth* of w in G and denote by $s(w, G)$ (or $s(w)$ for short). If iterational depth of w in G is finite, we call such iterated identity *bounded*.

We also say that G has *iterational depth* at most d if the depth of G with respect to any Engel type iterated identity that it satisfies, is at most d. The supremum of such d over all non-trivial (that is, not equivalent to the empty word in a free group) Engel type iterated identities w we denote by $s(G)$ and call it *iterational depth* of G. If G does not satisfy any non-trivial iterated identity, we put $s(G) = -\infty$.

defbounded

Definition 3.2. We say that G is bounded if $s(G)$ is finite (that is, $1 \leqslant s(G) < \infty$). If $s(G) = +\infty$ we say that G is fractal with respect to identities.

boundedandidentities

Remark 3.3. i) Any bounded group satisfies a non-trivial identity.

ii) If G is such that $s(G) < \infty$ then G is bounded if and only if G satisfies a non-trivial identity.

Proof. i) By definition, since G is bounded, there exists a non-trivial iterated identity w satisfied by G. Since G is bounded, it implies that there exists D such that for any $x_1, x_2, \dots, x_n$ it holds $w'(x_1, \dots, x_n) = e$ for at least one w' among w, $w_{\circ 2}$, …, $w_{\circ D}$. If w is not freely equivalent to the empty word, then none among w, $w_{\circ 2}$, …, $w_{\circ D}$ is freely equivalent to a free word (see i) of Example 2.4). If any tuple of elements of a group satisfy at least one among a finite list of identities, then the group satisifies a non-trivial identity.

examplefreesubgroup

ii) If G satisfies a non-trivial identity w, then G satisfies the (non-trivial) iterated identity w. Therefore, by definition of iterational depth $s(G) \neq -\infty$. Therefore, if $s(G) < \infty$, then $s(G)$ is a finite number, and thus G is bounded.

If G is bounded, then by the first part of the remark we know that G satisfies a non-trivial identity. Therefore, $s(G) \neq \infty$, and thus G is bounded so far as $s(G) \neq -\infty$.

□

examplefinite

Example 3.4. *Any finite group is bounded.*

Proof. Since G satisfies the E type iterated identity w, we know that for any $x_1, \dots, x_l$ in G there exists n such that $w_n(x_1, \dots, x_l) = e$. Put $s = \max n$, where maximum is taken over all possible choices of $x_1, \dots, x_l$ in G.

□

exampleabelian

Example 3.5. $s(\mathbb{Z}^d) = 1$. *Similarly, if G is an Abelian group which admits an element of infinite order, then $s(G) = 1$.*

Proof. Observe that if $w(x_1, \dots, x_n)$ is a E type iterated identity of G, then for any i the total number of x_i in w is equal to zero. (Indeed, otherwise we put $x_j = e$ for any $j \neq i$ and let x_i be an element of infinite order in G, we observe that $w_{\circ d}(x_1, x_2, \dots x_n) \neq e$ for any $d \geqslant 1$ and we get a contradiction). Since G is Abelian, we observe that for any $x_1, \dots, x_n \in G$ it holds $w(x_1, \dots, x_n) = 0$.

□

Remark 3.6. If G is a finitely generated Abelian group, then $s(G)$ is finite (indeed, in this case it is finite in case if G is finite by Example 3.4 and is equal to one in case if G is infinite by Example 3.5). exampleinfinite exampleabelian

It is clear that an infinitely generated Abelian group can have an infinite identity depth, as for example an infinite direct some $\sum_{i \geqslant 1} \mathbb{Z}/p^i\mathbb{Z}$ which satisfies an iterated identity $w(x_1) = x_1^p$ and does not satisfy any identity of the form $w'(x_1) = x_1^{p^k}$.

4. Fractal examples among elementary amenable groups

ectionfractalexamples
pleelementaryamenable

Example 4.1. *i) Let G be an extension of group of upper uni-triangular infinite square matricies with integer coefficients by a cyclic group (that acts on $\mathbb{Z} \times \mathbb{Z}$ by $z_1, z_2 \to z_1 + 1, z_2 + 1$). Consider $\bar{w}(x_1, x_2, x_3) = [x_1, [x_2, x_3]]$. Then G satisfies the iterated identity $\bar{w}$. For any iterated identity w of G it holds $s(w, G) = \infty$. In particular, G is fractal with respect to identities.*

ii) Moreover, there exists a continuum of non-isomorphic finitely generated elementary amenable groups that are fractal with respect to identities, and which do not satisfy any identity.

Proof. i) Denote by M_∞ the group of infinite upper uni-triangular matrices with integer coefficients (which we denote by $m_{i,j}$, $i, j \in Z$). This group is generated by upper triangular matrices m_i, $i \in \mathbb{Z}$ such that $m_i(i, i+1) = 1$ and $m_i(j, k) = 0$ if $j \neq k$ and $(j, k) \neq (i, i+1)$. We denote by ϕ the group homomorphism that sends m_i to m_{i+1}. The group G is by definition the extension of M_∞ by ϕ. It is clear that G is a finitely generated group, indeed, it is generated by ϕ and m_0. G is elementary amenable, since it is an extension of a locally nilpotent group M_∞ by a cyclic group.

lemmaiteridw

Lemma 4.2. *G satisfies the iterated identity $\bar{w} = \bar{w}(x_1, x_2, x_3) = [x_1, [x_2, x_3]]$.*

Proof. First observe that for any $x_2, x_3 \in G$ it holds $[x_2, x_3] \in [G, G] \subset M_\infty$ and for any $x_1 \in G$ it holds $\bar{w}(x_1, x_2, x_3) \in [G, G] \subset M_\infty$. Take some x_1, x_2, x_3 and consider $x = [x_2, x_3]$ and $y = \bar{w}(x_1, x_2, x_3)$. Since $x, y \in M_\infty$, they generate a subgroup which is nilpotent, and hence Engel. This shows that there exists $d \geqslant 1$ such that d-th commutator of the form $[\dots[[x, y], y] \dots, y]$ is equal to e. This implies that $\bar{w}_{\circ(d+1)}(x_1, x_2, x_3) = e$.

□

Now we want to show that for any iterated identity w of G it holds $s(w, G) = \infty$.

mmaMwithoutidentities

Lemma 4.3. *For any (non-iterated) non-trivial identity w there exists n such that the group $M^{(n)}$ of $n \times n$ upper uni-triangular matrices with integer coefficients does not satisfy w.*

Proof. Any torsion-free finitely generated nilpotent group can be imbedded as as subgroup in $M^{(n)}$ for some $n \geqslant 1$ (by a result of Jennings, see [20]). Let w be a word which is not mathtfnilpotentgroups

freely equivalent to the empty word. It is sufficient to check that there exists a torsion-free finitely generated nilpotent group which does not satisfy w. The latter follows from the fact that the free group F_2 does not satisfy w, and that free groups are residually torsion-free nilpotent ([21]).

□

Lemma 4.3 and Remark 3.3 imply that for any iterated identity w of G it holds $s(w, G) = \infty$. Since $\bar{w}$ is a non-trivial iterated identity of G (by Lemma 4.2), this implies, in particular, that G is fractal with respect to identities.

ii) Let G be the group constructed in the proof of i). Let G_1 be any finitely generated solvable group of step 3 such that the quotient over the the center $G/C(G) = \mathbb{Z} \wr \mathbb{Z}$. Consider the direct product $G' = G_1$. Observe that G_1, and hence G' satisfies the iterated identity w from the proof of i). Since G is a quotient of G', this implies that $s(w, G') \geqslant s(w, G) \geqslant \infty$, and hence $s(G') = \infty$, in other words G' is fractal with respect to identities. Observe also that $C(G)$ is trivial, and hence $C(G + G_1) = C(G)$. By [19], Section 3, we know that any countable Abelian group is a center of some group G_1 as above. Therefore, any countable Abelian A group there exists G', as above, with the center equal to A. In particular, there exists a continuum of such groups G_1,

Finally observe, that any G_1 as above does not satisfy a non-trivial identity, since G_1 contains G as a subgroup and since G does not satisfy any non-trivial identity.

□

The following example shows that , in contrast with usual identities, a conjugate of an iterated identity is not necessarily an iterated identity.

conjugatenotidentity

Example 4.4. *Let G and $\bar{w}$ be as in Example 4.1. Consider $w'(x_1, x_2, x_3, x_4) = x_4\bar{w}(x_1, x_2, x_3)x_4^{-1}$. Then G does not satisfy w'.*

Proof. Let $e_{i,j}(\lambda)$, $j > i$ denotes the elementary upper-diagonal unitary matrix m, such that $m_{i,j} = \lambda$ and all the other entries of m over diagonal are zero. Denote by $e_{i,j} = e_{i,j}(1)$.

Lemma 4.5. *Let α be an automorphism the group of infinite upper uni-triangular matrices M_∞ such that $\alpha(M)_{i,j} = M_{i+1,j+1}$ Given $y \in M_\infty$ consider the map $m_y : M_\infty \to M_\infty$ which sends x to $\alpha(xyx^{-1}y^{-1})$.*

If $y = e_{-1,0}e_{0,1}(-1)e_{-1,1}(-1)$ and $x = e_{-1,0}$, then $m_{y,\circ d}(x) \neq e$ for any $d \geqslant 1$. Moreover, $m_{y,\circ d}(x) \neq e$ is an elementary matrix with exactly one non-zero value off the main diagonal, at $(-d, 0)$.

Proof. First note that α preserves the lower central series $M = M_0 \supset M_1 \subset M_2 \ldots$ of M_∞ (here $M_{i+1} = [M, M_i]$), and therefore $m_{y,\circ d}(x) \in M_d$ for any $d \geqslant 1$. This implies that the entries on d diagonal above the main diagonal (that is, the entries at position$(k, k + l)$, $l \leqslant d$) of the matrix $m_{y,\circ d}(x)$ are zero.

Now prove by induction on d the fact that $m_{y,\circ d}(x) \neq e = e_{-(d+1),0}(\lambda)$, for some $\lambda \neq 0$.

The base $d = 0$ follows from the definition of x.

Suppose that $m_{y,\circ d}(x) = e_{-(d+1),0}(\lambda)$, for some $\lambda \neq 0$.

Observe that all non-zero entries of $[e_{-(d+1),0}(\lambda), y]$ off the main diagonal belong to the square $i \geqslant -(d + 1), i \leqslant 1$, $j \geqslant -(d + 1), j \leqslant 1$. Indeed, observe that if a matrix belongs to this square, then its inverse also belongs to this square, and if two matrices belong to this square, then their product belongs to this square. Observe also that $[e_{-d+1,0}(\lambda), y]$ belongs to M_{d+1}, and this is an uni-triangular matrix with $d + 1$ zero diagonals above the

main diagonal. We conclude that the only possible value of $[e_{-d+1,0}(\lambda), y]$ off the main diagonal at $i = -(d+1)$, $j = 1$. Let us show that this entry is non-zero. To do this observe that $e_{-d+1,0}$ does not commute with y. Indeed, the entry at $-(d+1,1)$ of $e_{-(d+1),0}(\lambda)y$ is equal to $-\lambda$, and that the the entry at $-(d+1,1)$ of $ye_{-(d+1),0}(\lambda)$ is equal to zero.

We have shown that $[e_{-d+1,0}(\lambda), y]$ is $e_{-(d+1,1)}(\lambda')$, for some $\lambda' \neq 0$. This implies that $\alpha([x,y]) = \alpha([e_{-(d+1),0}(\lambda), y]) = e_{-(d+2),0}(\lambda')$, and this concludes the proof of the induction step and the proof of the lemma.

□

Now take $x_2, x_3 \in G$ such that $[x_2, x_3] = y$. Such x_2 and x_3 do exist: indeed, take $x_2 = m_0$, $x_3 = \phi$, observe that $x_2 x_3 x_2^{-1} x_3^{-1} = m_0 \phi m_0 \phi^{-1} = m_0 m_1$. Put $x_1 = x = e_{-1,0}$ and $x_4 = \phi$.

Observe that for any $x \in M$ it holds $w'(x, x_2, x_3, x_4) = m_y(x)$. Therefore $w'_\circ(x_1, x_2, x_3, x_4) = m_{y,\circ d}(x_1) \neq e$ for any $d \geqslant 1$. We have therefore shown that w' is not an iterated identity of G.

□

5. Identities and iterated identities of group extensions, restriction of verbal map on an invariant subgroup and dynamics of its iterations

ectionnormalsubgroups

Let $w(x_1, \ldots, x_n)$ be a word and G be a group. Fix $x_2, \ldots, x_n$ and consider a verbal map from G to G, $x \to w(x, x_2, \ldots, x_n)$. We denote this verbal map by $\phi_w = \phi_{w,x_2,\ldots,x_n}$.

Now let N be a normal subgroup of G and suppose that w is an identity of G/N. Then for any $x_2, \ldots, x_n \in G$ and any $x \in G$ it holds $w(x, x_2, \ldots, x_n) \in N$. In other words $\phi_{w,x_2,\ldots,x_n}(x) \in N$ for any $x \in G$, and, in particular, for any $x \in N$. We can therefore study the restriction of the verbal map $\phi_{w,x_2,\ldots,x_n}$ to N.

remarkuv **Remark 5.1.** Observe than any word $w = w(x_1, \ldots, x_n)$ can be written as $w(x_1, \ldots, x_n) = u(x_1, \ldots, x_n)v(x_2, \ldots, x_n)$, where the word u has the following form

$$u(x_1, \ldots, x_n) = \prod_{i=1}^{N} \alpha_i x_1^{l_i} \alpha_i^{-1},$$

where $l_i \in \mathbb{Z}$ and α_i are some word in $x_2, \ldots, x_n$.

If w is an identity of G/N, then so is $v(x_2, \ldots, x_n)$. In other words $v(x_2, \ldots, x_n) \in N$ for any $x_2, \ldots, x_n \in G$

kernelinN **Remark 5.2.** Let $w(x_1, \ldots, x_n)$ be a word, G be a group and N is a normal subgroup of G. Fix some $x_2, \ldots, x_n$ and consider the verbal map $\phi = \phi_{w,x_2,\ldots,x_n}$ (that is, $\phi(x) = w(x, x_2, \ldots, x_n)$. Then either $\phi(n) \in N$ for any $n \in N$, or $\phi(n) \notin N$ for any $n \in N$.

remarkuv

Proof. Use Remark 5.1 and write $w = u(x_1, \ldots, x_n)v(x_2, \ldots, x_n)$. Since u is a product of conjugates of x_1 and since N is a normal subgroup, we conclude that $\phi_u(x) \in N$ for any $x \in N$. If $v_{x_2,\ldots,x_n} \in N$, then $\phi(x) \in N$ for any $x \in N$.

Otherwise, if $v_{x_2,\ldots,x_n} \notin N$, then $\phi(x) \notin N$ for any $x \in N$.

□

remarkuv

lemmahomom **Lemma 5.3.** *Consider w, v and u defined in Remark 5.1. Fix $x_2, \ldots, x_n$. Consider the verbal map ϕ_u from G to G and its restriction to N.*

If N is an Abelian normal subgroup, then the restriction of ϕ_u to N defines a group homomorphism on N.

Proof. Let $y, z \in N$. Observe that $(yz)^{l_i} = y^{l_i} z^{l_i}$, and hence

$$\phi_u(yz) = \prod_{i=1}^{N} \alpha_i (yz)^{l_i} \alpha_i^{-1} = \prod_{i=1}^{N} \left(\alpha_i y^{l_i} \alpha_i^{-1} \alpha_i z^{l_i} \alpha_i^{-1}\right) = \prod_{i=1}^{N} \alpha_i y^{l_i} \alpha_i^{-1} \prod_{i=1}^{N} \alpha_i z^{l_i} \alpha_i^{-1} = \phi_u(y)\phi_u(z).$$

□

commutewithaction

Lemma 5.4. . *We consider again w, u, v from Remark 5.1. Let G/N and N be Abelian groups. Then*

i) for any $x_2, \dots, x_n \in G$ the restriction of ϕ_u to N commutes with the action (by conjugation) of G on N.

ii) for any $x_2, \dots, x_n \in G$ any any $x \in N$ $\phi_u(x)$ belongs to the linear span of $a(x)$, $a \in G/N$.

Proof. First observe that the action of g on N depends only on the image of g in G/N. Indeed, if $g = hn$, $n \in N$, then $gmg^{-1} = hnmn^{-1}h^{-1} = hmh^{-1}$ (n commute with m since $m, n \in N$ and N is Abelian). To prove i) observe that by the definition of ϕ_i in Remark 5.1 $\phi_u(n) = \prod \alpha_i n^{l_i} \alpha_i^{-1}$, and hence

$$g\phi_u(n)g^{-1} = g \prod \alpha_i n^{l_i} \alpha_i^{-1} g^{-1} = \prod g\alpha_i n^{l_i} \alpha_i^{-1} g^{-1} = \prod g\alpha_i n^{l_i} (g\alpha_i)^{-1}$$

Since G/N is Abelian, $g\alpha_i = \alpha_i g$ in G/N, and hence the action by conjugation on N of these two elements are the same. Therefore

$$g\phi_u(n)g^{-1} = \prod \alpha_i g n^{l_i} (\alpha_i g)^{-1} = \prod \alpha_i (gng^{-1})^{l_i} \alpha_i^{-1} = \phi_u(gng^{-1}).$$

ii) follows from the definition of ϕ_u. □

Two following lemmas control the depth of a group which contains a cyclic normal subgroup, finite or infinite.

afinitenormalsubgroup

Lemma 5.5. *Let G be a group and N be a finite normal subgroup of G. The cardinality of N we denote by $\#N$. Let w be an iterated identity of G. Then*

$$s(w, G) \leqslant (s(w, G/N) + 1)(\#N + 1)$$

In particular, If G/N is bounded, then G is bounded.

Proof. Ift $w(x_1, \dots, x_n)$ be an iterated identity of G, then w is an iterated identity of G/N. Denote by D the iterational depth $s(w, G/H)$. For any $x_1 \dots, x_n \in G/N$ there exists $d \leqslant D$ such that $w_{\circ d}(x_1, \dots, x_n) = e$ in G/N. In other words, for any $x_1, \dots, x_n$ in G here exists $d \leqslant D$ such that $w_{\circ d}(x_1, \dots, x_n) \subset N$. Fix $x_2, \dots, x_n$ and consider some $x \in N$. Since w is an iterated identity of G there exists m (depending on x and $x_2, \dots, x_n$) such that $w_{\circ m}(x, x_2, \dots, x_n) = e$.

Let $\phi : G \to G$ be the verbal map $\phi(x) = w(x, x_2, \dots, x_n)$. Let R denotes the cardinality of N. Observe that for any $x \in G$ there exist $1 \leqslant m_1 < m_2 < m_3 < m_{R+1}$ such that $\phi_{\circ m_j}(x) \in N$ for any $j : 1 \leqslant j \leqslant R + 1$ and such that $m_1 \leqslant D + 1$, such that $m_{j+1} - m_j \leqslant D + 1$ for any $j \leqslant R$ and such that

Observe that $1 \leqslant m_1 < m_2 < m_3 < m_{R+1} \leqslant (D + 1)(R + 1)$. Therefore, there exist at least among these $R + 1$ elements there exist at least two equal ones, that is, there exist $r, s : 1 \leqslant r < s \leqslant (R + 1)(D + 1) =$ such that $\phi_{\circ r}(x) = \phi_{\circ s}(x)$, This implies that for any $t \geqslant 0$ $\phi_{\circ r+t}(x) = \phi_{\circ s+t}(x)$. This implies that if $\phi_{\circ m}(x) = e$ for some $m \geqslant 1$, then there exists $m \leqslant (R + 1)(D + 1)$ such that $\phi_{\circ m}(x) = e$

We have shown therefore that $s(w, G) \leqslant (s(w, G/N) + 1)(\#N + 1)$.

Now assume that G/N is bounded. Then $s(G/N) < \infty$, and thus $s_G < \infty$. Since G/N is bounded, we know from Remark 3.3 that G/N satisfies at least one non-trivial identity, which we denote by w. Then G satisfies an identity w^R, and it is clear that w^R is not freely equivalent to a trivial word. Thus, we know that $s(G) < \infty$ and that G satisfies a non-trivial identity. Using 3.3 we conclude that G is bounded.

□

lemmacyclicquotient

Lemma 5.6. *Let G be a group and N be an infinite normal cyclic subgroup of G. For any iterated identity w of G it holds $s(w, G) \leqslant 2s(w, G/N) + 1$.*

In particular, if If G/N is bounded, then G is bounded.

Proof. Let $w(x_1, \dots, x_n)$ be an iterated identity of G. Since w is an iterated identity of G/N and since G/N is bounded, there exists D such that for any $x_1, \dots, x_n$ in G here exists $d \leqslant D$ such that $w_{\circ d}(x_1, \dots, x_n) \subset N$.

Fix $x_2, \dots, x_n$. For any $d \geqslant 0$ consider the verbal map ϕ_d, defined by $w_{\circ d}$. By remark 5.2 we know that either $\phi_d(x) \in N$ for any $x \in N$, or $\psi_d(x) \notin N$ for any $x \in N$. This implies that if $d'' < d'$ and $\phi_d(e), \phi''_d(e) \in N$, then $\phi_{d'-d''}(e) \in N$. Let D be a common divisor of such d'. We have $\phi_l(e) \in N$ if and only if l is divided by D, and , moreover, for any $x \in N$ we know that $\phi_l(e) \in N$ if and only if l is divided by D. In particular, if $\phi_l(x) = e$ for some $x \in N$, then l is divided by D. Observe that $D \leqslant d + 1$, since there exists $m : 1 \leqslant m \leqslant d + 1$ such that $\phi_m(e) \in N$.

Apply Remark 5.1 to $w_{\circ D}$ and consider u and v defined in this remark. For any $x_2, \dots, x_n \in G$, the restriction of the verbal map of $w_{\circ D}$ has the form $\psi(x) = \phi_u(x) + v$, where ϕ is a homomorphism of (an infinite cyclic group) N, and $v \in N$.

We have already shown that for any $x \in N$ there exists an integer l such that $\phi_{Dl}(x) = \psi_{\circ l}(x) = e$.

Since ϕ_u is a homomorphism from $\mathbb{Z}$ to $\mathbb{Z}$, there exists an integer T such that $\phi_u(x) = Tx$. Since $\psi_{\circ m(e)} = 0$ for some m, it follows that $v + \phi_u(v) + \dots \phi_{u,\circ(m-1)}(v) = 0$. This implies that $\phi_u(e)$ takes finitely many values, we denote this finite set by Ω. Observe that for any $l \geqslant 1$ there exists $\omega \in \Omega$ such that $\psi_{\circ l}(x) = T^l x + \omega$, for any $x \in N$. Assume that $T \neq 0$. Note that then for any sufficiently large x all values of $\psi_{\circ l}(x)$ are non-zero. This contradiction shows that $T = 0$.

We know , therefore, that $\phi_(u)(x) = 0$ for any $x \in N$, and hence $\psi_{\circ l}(x) = v$ for any $l \geqslant 1$, and hence $v = e$. This shows that $\psi(x) = 0$ for any $x \in N$, and this implies that for any $x_1, \dots, x_n$ there exists $l \leqslant d + D \leqslant 2d + 1$ such that $w_{\circ l}(x_1, \dots, x_n) = e$. We have shown therefore that $s(w, G) \leqslant 2s(w, G/N) + 1$.

Finally, observe that G satisfies a non-trivial identiy if G/N satisfies a non-trivial identity. Indeed, let $w(x_1, \dots, x_n)$ be an identity of G/N. Consider $w'(x_1, \dots, x_{2n}) = [w * (x_1, \dots, x_n), w(x_{n+1}, \dots, x_{2n})]$. It is clear that w' is not freely equivalent to an empty word if w has this property, and it is clear that G satisfies the identity w'. □

Lemma 5.5 and Lemma 5.6 imply the following

Corollary 5.7. *Any finitely generated polycyclic group is bounded.*

5.1. **Nilpotent groups.**

remarkuvnilpotent

Remark 5.8. Any word $w(x_1, \dots, x_n)$ can be written as

$$w(x_1, x_2, \dots, x_n) = \prod_{j=1}^{N} [x_1^{l^j}, u_j] x_1^r v(x_2, \dots, x_n),$$

where u_j are words in $x_2, \dots, x_n$, l_j are integers and r is the total number of x_1 in w . We denote

$$u(x_1, \dots, x_n) = \prod_{j=1}^{N} [x_1^{l^j}, u_j].$$

Proof. Write a word w as a product of words $(x_2, x_3, \dots, x_n)$ and powers of x_1 and prove the statement of the remark by induction on the number of terms in such product.

□

It is not difficult to describe all E type iterated identities of a nilpotent group:

examplenilpotent

Example 5.9. *Let G be a nilpotent group and $w(x_1, x_2, \dots, x_n)$ be a word. Let u, v be the words defined in Remark 5.8 and r be the total number of x_1 in w.*

remarkavnilpotent

i) Suppose that there exists an integer $m > 0$ such that for any element $g \in G$ there exists an integer $t > 0$ such that such that $g^{m^t} = e$. Take m to be the minimal positive integer with this property. The word w is an E type iterated identity of G if and only if r is divided by m and $v(x_2, \dots, x_n)$ is an identity of G.

ii) Otherwise (for example, if G contains at least one element of infinite order) the word w is an E type iterated identity of G if and only if $r = 0$ and $v(x_2, \dots, x_n)$ is an identity of G.

Proof. First suppose that w is an E type iterated identity of G. Consider $x_1 = x$, $x_2 = x_3 = \dots x_n = e$. Observe that for all $d \geqslant 1$ it holds $w_{\circ d}(x_1, x_2, \dots, x_n) = x^{r^d}$. This implies $r = 0$ that under assumption of ii) and that r is divided by m under assumption of i).

Let us show that v is an identity of G. Suppose not. Then there exists $x_2, \dots, x_n$ such that $v(x_2, \dots, x_n) \neq e$.

If G satisfies the assumption of ii), take $M \geqslant 0$ such that $v(x_2, \dots, x_n)$ belongs to the M-th term of the lower central series of G and does not belong to the $M+1$-th term of this series. Observe that for any x_1 and any $d \leqslant M$ the iteration $w_{\circ d}(x_1, x_2, \dots, x_n)$ belongs to the d-th term of the lower central series of G. Observe also that $w_{\circ (M+1)}(x_1, x_2, \dots, x_n)$ (as well as $w_{\circ d}(x_1, x_2, \dots, x_n)$ for any $d \geqslant M+1$) belongs to the M-th term of the lower central series but does not belongs to the $M+1$-th term. This shows that $w_{\circ d}(x_1, x_2, \dots, x_n)$ is not equal to e, for any x_1 and any $d \geqslant M$. This is a contradiction with the fact that w is an iterated identity of G.

If G satisfies the assumption of i), observe that any finitely generated subgroup of G is finite. Consider the following central series of G: $\lambda_0(G) = G$, $\lambda_{i+1}(G)$ be the normal subgroup generated by $[\lambda_i(G), G]$ and $\lambda_i(G)^m$. We consider, as before, $x_2, \dots, x_n$ such that $v(x_2, \dots, x_n) \neq e$. Put $x_1 = e$. Let G' be the subgroup of G generated by $x_1, x_2, \dots, x_n$. Observe that the series $\lambda_d(G')$ (of the finite group G') has a finite number of non-trivial terms. Since $v(x_2, \dots, x_n) \neq e$, there exists $M \geqslant 0$ such that $v(x_2, \dots, x_n) \neq e$ belongs to $\lambda_M(G')$ and does not belong to $\lambda_{(M+1)}(G')$.

Observe that for any x_1 and any $d \leqslant M$ $w_{\circ d}(x_1, x_2, \dots, x_n)$ belongs to $\lambda_d(G')$. Observe also that $w_{\circ M+1}(x_1, x_2, \dots, x_n)$ (as well as $w_{\circ d}(x_1, x_2, \dots, x_n)$ for any $d \geqslant M+1$) belongs to $\lambda_M(G')$-th but does not belongs to $\lambda_{M+1}(G')$-th term. This contradicts the fact that w is an iterated identity of G.

Now suppose that r_i satisfy the assumption of i) and ii) correspondingly. Let us show that w is an E type iterated identity of G.

Under assumption of ii) it is clear that $w_{\circ d}(x_1, x_2, \dots, x_n) = x^N$ belongs to the d-the term of the lower central series of G.

If G satisfies the assumption of i) and w satisfies the condition of i), it is clear that $w_{\circ d}(x_1, \dots, x_n)$ belongs to $\lambda_d(G')$, where G' is a subgroup generated by $x_1, \dots, x_n$. Observe that G' is a finite nilpotent group such that for any $g \in G'$ there exists t such that $g^{m^t} = e$. This implies that $\lambda_d(G')$ has finite length (that is, there exists a finite number of non-zero terms in this series), and therefore w is an iterated identity of G.

□

Corollary 5.10. *Let G be a nilpotent group. Let $\Omega_n \subset F_n$ be the set of E type iterated identities of G. Then Ω_n is a normal subgroup of the free group F_n, for any $n \geqslant 1$.*

Proof. If G satisfies the assumption of ii) and w_1 and w_2 satisfy the conditions of ii), observe that their product $w = w_1 w_2$, as well as for any conjugate $w = w' w_1 (w')^{-1}$ it holds $w_{\circ d}(x_1, \dots, x_n) = u_{\circ d}(x_1, \dots, x_n)$ belongs to the d-th term of the lower central series of G.

Observe also that if G satisfies the assumption of i) and w_1 and w_2 satisfy the conditions of i), then their product $w = w_1 w_2$, as well as any conjugate $w = w' w_1 (w')^{-1}$ it holds $w_{\circ d}(x_1, \dots, x_n)$ belongs to $\lambda_d(G')$, where G' is a (finite nilpotent) subgroup generated by $x_1, \dots, x_n$.

□

6. Examples of iterational depth. Wreath products

sectionwreath

propositionwithz

Proposition 6.1. $s(\mathbb{Z} \wr \mathbb{Z}) = 2$

Proof. Let $G = \mathbb{Z} \wr \mathbb{Z}$, $N = \mathbb{Z}^\infty \subset G$. Let $A = G/N$. We know that $A = \mathbb{Z}$. Let $w(x_1, \dots, x_n)$ be an iterated E-type identity of G. Since G has elements of infinite order, the total number of x_i in w is equal to zero, for each i. Therefore, w is an identity of A. We consider $u(x_1, x_2, \dots, x_n)$ and $v(x_2, \dots, x_n)$ defined in Remark 5.1. It holds $w(x_1, \dots, x_n) = u(x_1, x_2 \dots x_n) v(x_2, x_3, \dots x_n)$.

remarku

We want to show that for any $x \in N$ it holds $\phi_u(x) = 0$. Let ϵ_i, $i \in \mathbb{Z}$ be the standard basis of $\mathbb{Z}^\mathbb{Z}$. If $\phi_u(x) \neq 0$ for some $x \in N$, then there exists $i \in \mathbb{Z}$ such that $\phi(\epsilon_i) \neq 0$. Let

$$\phi_u(\epsilon_i) = \sum_{i=-M}^{M} \beta_j \epsilon_{i+j}$$

commutewithaction

By Lemma 5.4 we know that ϕ_u commutes with the action of $\mathbb{Z}$ on $\mathbb{Z}^\mathbb{Z}$ by conjugation. Therefore, for any $k \in \mathbb{Z}$ it holds

$$\phi_u(\epsilon_k) = \sum_{i=-M}^{M} \beta_j \epsilon_{k+j}$$

Let m be the maximal among i such that $\beta_i \neq 0$ and suppose that $m \geqslant 0$. Let

$$v = v(x_2, x_3, \dots, x_n) = \sum_{i=-L}^{L} \gamma_i \epsilon_i$$

Take any $k > L$. Observe that the l-the iteration of $\phi_u v$, applied to ϵ_k, has a coordinate $\beta_m^l \neq 0$ at the basis vector ϵ_{k+ml}. Therefore, any such iteration is not equal to zero, and hence no E type iteration of w at $(x_1, \dots, x_n)$ is equal to e in G. We have got a contradiction with the fact that w is an iterated identity of G.

Therefore, for any $x_2, \dots, x_n$ and any $x \in N$ it holds $\phi_u(x) = 0$. Take any $x_1 \in G$. Since $w(x_1, \dots, x_n \in N$, we conclude that for any $l \geqslant 2$ the l-the iteration of w at

$x_1, \dots, x_n$ is equal to $v(x_2, \dots, x_n)$. Since for some l this iteration is trivial, we see that $v(x_2, \dots, x_n) = e$ and hence the second iteration of w is trivial.

We have shown that $s(\mathbb{Z} \wr Z) \leqslant 2$.

Remark 6.2. Take $w = (x_1, x_2) = [x_1, [x_1, x_2]]$.

1) w is an iterated identity of $\mathbb{Z} \wr \mathbb{Z}$.

2) w is not an identity of $\mathbb{Z} \wr \mathbb{Z}$.

Proof of the remark.

1) Note that $w = (x_1, x_2) = [x_1, [x_1, x_2]]$ belongs to the commutator subgroup of $\mathbb{Z} \wr \mathbb{Z}$ for any x_1 and x_2. Observe also that for any x_1, x_2 their commutator belongs to the commutator subgroup, and the commutator subgroup of $\mathbb{Z} \wr \mathbb{Z}$ is Abelian. Therefore, $w(x_1, x_2) = e$ for any $x_2 \in \mathbb{Z} \wr \mathbb{Z}\, x_1$ and any x_1 in the commutators subgroup of $\mathbb{Z} \wr \mathbb{Z}$. Hence, the second E type iteration of w is trivial.

2) Let x, a be the standard generators of $\mathbb{Z} \wr \mathbb{Z}$, where x corresponds to the generator of $\mathbb{Z}$ that acts. Put $x_1 = x$, $x_2 = xa$. Observe that $[x_1, x_2]$ is not a power of x, and this implies that $[x_1, x_2]$ does not commute with x. In other words, $w(x_1, x_2) \neq e$.

qed

The remark implies that $s(\mathbb{Z} \wr \mathbb{Z}) \geqslant 2$, and hence $s(\mathbb{Z} \wr \mathbb{Z}) = 2$

□

roppositionzwithfinite **Proposition 6.3.** *$s(\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}) = k + 1$, k being the maximum of α_i, where $R = \prod p_i^{\alpha_i}$, p_i are prime numbers, $\alpha_i \in \mathbb{Z}_+$.*

Proof. We have $G = \mathbb{Z} \wr \mathbb{Z}/n\mathbb{Z}$ and we consider $N = (\mathbb{Z}/R\mathbb{Z})^{\infty} \subset G$. $A = G/N = \mathbb{Z}$. Let w be an iterated identity of G. Since G has elements of infinite order, the total number of x_i in w is equal to zero, for each i. Therefore, w is an identity of A. We consider $u(x_1, x_2, \dots, x_n)$ and $v(x_2, \dots, x_n)$ defined in Remark 5.1. It holds $w(x_1, \dots, x_n) = u(x_1, x_2 \dots x_n)v(x_2, x_3, \dots x_n)$.

remarkuv

We want to show that for any $x \in N$ it holds $\phi_{u,\circ k}(x) = 0$. Let ϵ_i, $i \in \mathbb{Z}$ be the standard basis of $(\mathbb{Z}/R\mathbb{Z})^{\mathbb{Z}}$.

Suppose that $\phi_{u,\circ k}(x) \neq 0$ for some $x \in N$. Since ϕ_u, and hence also all its iterations $\phi_{u,\circ k}$ are homomorphism of the Abelian group N, exists $i \in \mathbb{Z}$ such that $\phi_{u,\circ k}(\epsilon_i) \neq 0$. Let

$$\phi_u(\epsilon_i) = \sum_{i=-M}^{M} \beta_j \epsilon_{i+j}$$

commutewithaction By Lemma 5.4 we know that ϕ_u commutes with the action of $\mathbb{Z}$ on $(\mathbb{Z}/R\mathbb{Z})^{\mathbb{Z}}$ by conjugation. Therefore, for any $r \in \mathbb{Z}$ it holds

$$\phi_u(\epsilon_r) = \sum_{i=-M}^{M} \beta_j \epsilon_{r+j}$$

Observe that if all b_i are divided by any of p_j among the prime divisors of R, then all coefficients of $\phi_{u,\circ k}(\epsilon_i)$ are divided by p_j^k, for all j. In this case all such coefficients are divided by R, and thus $\phi_{u,\circ k}(\epsilon_i) = 0$ in $(\mathbb{Z}/R\mathbb{Z})^{\mathbb{Z}}$.

Therefore, we can assume that there exists some p among prime divisors of R and some integer j such that β_j is not divided by p. Let m be the maximal among j such that β_j is not divided by p. Without loss of generality we can assume that $m \geqslant 0$.

Let

$$v = v(x_2, x_3, \dots, x_n) = \sum_{i=-L}^{L} \gamma_i \epsilon_i$$

Take any $T > L$. Observe that the l-the iteration of $\phi_u v$, applied to ϵ_M, has a coordinate $\beta_m^l \neq 0$ at the basis vector ϵ_{T+ml}. Therefore, any such iteration is not equal to zero, and hence no E type iteration of w at $(x_1, \dots, x_n)$ is equal to e in G. We have got a contradiction with the fact that w is an iterated identtity of G.

Now let us show that γ_i is divided by R, for any i. Indeed, otherwise there exist a prime divisor p of R such that R is divided by p^s, and there exists i such that γ_i is not divided by p^s. Let t be the maximal number such that γ_i is divided by p^t for all i. We know that $t < s$. Observe that all coefficients of $\phi_u, (v)$ are divisible by p^{t+1}, and hence $\phi_w, (v) \equiv v$ modulo $p^{t+1}(\mathbb{Z}/R\mathbb{Z})^{\mathbb{Z}}$. This implies that for all $l \geqslant 1$ $\phi_{w,\circ l}, (v) \equiv v$ modulo $p^{t+1}\mathbb{Z}/R\mathbb{Z}$. Therefore, $\phi_{w,\circ l} \neq e$ for all $l \geqslant 1$. This contradiction shows that γ_i is divided by R, for any i. In other words, $v \equiv 0$ in $(\mathbb{Z}/R\mathbb{Z})^{\mathbb{Z}}$.

We have shown therefore that for any x_2, x_3, ..., $x_n \in G$ and any $x \in N$ it holds $\phi_{w,\circ k}(x) = w_{\circ k}(x, x_2, \dots, x_n) = e$ This implies that $s(\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}) \leqslant k + 1$.

Remark 6.4. Take $w(x_1, x_2) = [x_1^m, x_2]$, where $m = p_1 \dots p_l$, p_i are prime numbers. Consider $G = \mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}$

1) If prime divisors of R are contained among $p_1, \dots, p_l$, then $w(x_1, x_2)$ is an iterated identity of $\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}$

2) If n is divided by p_1^k, then $w_{\circ k}(x_1, x_2)$ is not an identity of $\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}$.

Proof of the remark.

1) Observe that $w(x_1, x_2) \in (\mathbb{Z}/R\mathbb{Z})^{\infty}$ for any $x_1, x_2 \in G$ Observe also that for any $x_2 \in G$ and any $k \geqslant 0$ and $x_1 \in m^k(\mathbb{Z}/R\mathbb{Z})^{\infty}$ it holds $w(x_1, x_2) \in m^{k+1}(\mathbb{Z}/n\mathbb{Z})^{\infty}$. By the assumption of the remark we know that there exist k such that m^k is divided by R. Hence for any $x_1, x_2 \in G$ the $k+1$-th Engel type iteration of w, evaluated at x_1, x_2 is trivial.

2) Let x, a be the standard generators of $\mathbb{Z} \wr \mathbb{Z}R\mathbb{Z}$, where x corresponds to the generator of $\mathbb{Z}$. Put $x_1 = xa$, $x_2 = x$. Observe that $[x_1^m, x_2]$ is a configuration of the wreath product that has two non-zero values: -1 at $m+1$ and $+1$ at 1. Applying w to $[x_1^m, x_2]$ and x_2 we get a configuration, with the value at the maximal point of the support equal to m. The k-the iteration, applied to x_1 and x_2, has value m^{k-1} at one of the points of its support. Since R is divided by p_1^k, we know that $m^{k-1} \neq 0$ in $\mathbb{Z}/R\mathbb{Z}$. We conclude that k-iteration of w is not an identity of $\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}$.

qed

The remark implies that $s(\mathbb{Z} \wr \mathbb{Z}/R\mathbb{Z}) \geqslant k + 1$ for R, k satisfying the assumption of the theorem).

□

6.1. **Central extensions.**

kernelnormalsubgroup

Remark 6.5. Let H be a normal subgroup of G, $w(x_1, \dots, x_n)$ be some word and denote by ϕ_d the verbal map $\phi_d(x) = w_{\circ d}(x, x_2, \dots, x_n)$.

i) For any $x_2, \dots, x_n$ there exists D (an integer number or ∞) such that the following holds. For any $x \in H$ the image $\phi_l(x) \in N$ if and only if l is divided by D.

ii) If we assume additionally that w is an iterated identity of G/H, then $D \leqslant s(w, G/H) + 1$.

Proof. i) Apply Remark 5.1 to $w_{\circ d}$. We know that there exist words u_d and v_d such that $\phi_d(x) = \phi_{u_d}(x, x_2, \dots, x_n)v_d(x_2, \dots, x_n)$. Observe that $\phi_{u_d}(x, x_2, \dots, x_n)$ is a product of conjugates of powers of x. Therefore $\phi_{u_d}(x, x_2, \dots, x_n)$ belongs to H whenever $x \in H$, since H is a normal subgroup. If $v_d(x_2, \dots, x_n)$ belongs to H then $\phi_d(x) \in H$ for any $x \in H$. If $v_d(x_2, \dots, x_n)$ does not belong to H then $\phi_d(x) \notin H$ for any $x \in H$.

Now observe that if D is such that $\phi_{\circ d}(x) \in N$ for any $x \in N$, then $\phi_{\circ kD}$ also has property. Observe also that if $a > b$ and $\phi_{\circ a}$ and ϕ_{circb} both have this property, then $\phi_{\circ(a-b)}$ also has this property.

ii) If $\phi_1(x) \in H$ for some $x \in H$, then D (defined in i)) is equal to 1, and thus $1 = D \leqslant s(w, G/H)$. Otherwise, $\phi_1(e) \notin H$. In this case there exists $l \leqslant s(w, G/H)$ such that $\phi_{l+1} = \phi_l(\phi(e)) \in H$, and thus $D \leqslant s(w, G/H) + 1$.

□

Lemma 6.6. *Assume that H is a central subgroup of G. For any iterated identity w of G such that the total number of each x_i in w is equal to zero*

$$s(w, G) \geqslant 2s(w, G/H) + 1.$$

In particular, if G contains at least one element of infinite order and if G/H is bounded, then G is bounded.

Proof. Let $d = s(w, G/H)$. Fix some $x_2, \dots x_n \in G$.

Let ϕ_d is a verbal map $\phi_d(x) = w_{\circ d}(x, x_2, \dots, x_n)$.

By Remark 6.5 we know that there exists $D \leqslant d + 1$ such that for any $x \in N$ the image $\phi_l(x) \in N$ if and only if l is divided by D.

Apply Remark 5.1 to $w_{\circ D}$ and write $\phi_D(x) = u(x, \dots, x_n)v(x_2, \dots, x_n)$. Since the total number of entries of x_1 in w is equal to zero and since H is central, we see that $u(x, \dots, x_n) = e$. Let us show that $v(x_2, \dots, x_n) = e$. Indeed, otherwise $w_{\circ(lD)}(e, x_2, \dots, x_n) = v(x_2, \dots, x_n) \neq r$ for any positive integer l. This implies that $w_{\circ m}(e, x_2, \dots, x_n) \neq e$ for any positive integer m, and this is a contradiction with the fact that w is an iterated identity of G.

We have shown therefore that $v(x_2, \dots, x_n) = e$, and thus $w_{\circ D}(x, x_2, \dots, x_n) = e$ for any $x \in H$.

This implies that for any $x_2, x_3, \dots, x_n$ and any $x \in G$ there exists $l \leqslant d + D \leqslant 2d + 1$ such that $w \circ l(x, x_2, \dots, x_n) = e$, and hence $s(w, G) \geqslant 2s(w, G/H) + 1$.

Finally observe that if G/H satisfies some non-trivial identity, so does G: if G satisfies an E type iterated identity $w(x_1, \dots, x_n)$, then G/H satisfies the commutator identity $w'(x_1, \dots, x_{2n}) = [w(x_1, \dots, x_n), w(x_{n+1}, \dots, x_{2n})]$. □

Example 6.7. *Let G be a solvable group such that the center $C(G) = \sum_{i=1}^{\infty} \mathbb{Z}/p^i\mathbb{Z}$ and such that $G/C(G) = \mathbb{Z} \wr \mathbb{Z}$. (Such groups do exist, see Section 3 in* [19]*). Then G is bounded,, but $C(G)$ is not bounded.*

Proof. From Proposition 6.1 we know that $\mathbb{Z} \wr \mathbb{Z}$ is bounded. Since G is a central extension of $\mathbb{Z} \wr \mathbb{Z}$, we can use Lemma 6.6 to conclude that G is bounded.

□

7. METABELIAN GROUPS

Theorem 7.1. *If G is a finitely generated metabelian group, then $s(G)$ if finite.*

Proof. If G is finite, we know that G is bounded (see Example 3.4). We will therefore assume that G is infinite.

Let N be an Abelian normal subgroup of G, such that $A = G/N$ is Abelian.

Let w be an iterated E-type identity of G. Since G is infinite finitely generated metabelian group, it contains an element of infinite order. Therefore, for each i the total number of x_i in w is equal to zero. In particular, w is an identity of A. Using Remark 5.1 we can write $w(x_1, \dots, x_n) = u(x_1, \dots, x_n)v(x_2, \dots, x_n)$.

Fix $x_2, \dots, x_n$ and consider the homomorphism $\phi_u : N \to N$. (By Lemma 5.3 we know that ϕ_u is indeed a homomorphism). We also consider $\phi_w(x) = \phi_u(x)v(x_2, \dots, x_n) = w(x, x_2, \dots, x_n)$.

Since A is a finitely generated Abelian group, $A = \mathbb{Z}^d + A_f$, $d \geqslant 0$, A_f is a finite Abelian group.

Given a map ρ from N to N let us say that ρ is recurrent if for any $x \in N$ there exists n such that the n-th iteration $\rho_{\circ n}(x) = e$. Since w is an iterated identity, we know that the map $\phi_w : N \to N$ is recurrent. Our goal is to find a common upper bound for (minimal) n such that $\phi_{w,\circ n}(x) = e$, which does not depend on $x_2, \dots, x_n$ and does not depend on w. To do this, we need to study dynamics of the homomorphism ϕ_u and of the "affine map" ϕ_w on N.

lemmafinitetensor

Lemma 7.2. *Let F be a torsion-free Abelian group such that its tensor product $F \otimes \mathbb{Q}$ has finite dimension, which we denote by D. $\xi : F \to F$ be a homomorphism, a is some element of F and $\rho(x) = \xi(x) + a$. Suppose that for any $x \in F$ there exists n such that $\rho_{\circ n}(x) = e$. Then for any $x \in F$ it holds $\rho_{\circ D}(x) = \xi_{\circ D}(x) = e$.*

Proof. Since F is torsion free, F is imbedded in $V = F \otimes \mathbb{Q}$.

It is clear that ξ extends to a linear map from V to V, and we use the same letter ξ to denote this linear map. First let us show that the kernel of $\xi_{\circ D}$ is equal to V. Let $V_i = \ker \xi_{\circ i}$. It is clear that $V_1 \subset V_2 \subset V_3 \dots$. It is also clear that if for some j it holds $V_j = V_{j+1}$, then for any $k \geqslant j$ it holds $V_k = V_j$. Suppose that the kernel of $\xi_{\circ d}$ is not equal to V. Then there exists a linear subspace U of dimension $\leqslant D - 1$ such that for any $i \geqslant D - 1$ it holds $V_i = U$.

Now observe that there exits M such that $\phi_{\circ M}(0) = \phi_{\circ(M-1)}(\phi(0)) = 0$. Note that ρ (as well as its iterations) extends to an affine map from V to V, denote by W_i the kernel of $\phi_{\circ i}$ in V.

Observe that that W_i is contained in the union of the image of U by a finite number of translations. In other words, the union $W = \cup W_i$ is contained in $\cup_{i=1}^{l}(U + v_i)$, for some $l \geqslant 0$ and some $v_i \in V$. Note that since the linear span of the image of F is equal to $V \neq U$, there exists a $f \in F$ such that its image does not belong to U. Replacing, if necessary, f by some of its power Lf, we conclude that there exists $f \in F$ such that its image in the tensor product does not belong to W. This is a contradiction with the assumption of the lemma.

Now we know that the kernel of $\xi_{\circ D}$ is equal to V: $V_D = V_{D+1} = V_{D+2} = \dots = 0$ and $\xi_{\circ D}(x) = 0$ for any $x \in F$.

Observe that $\phi_{\circ j}(0) = a + \xi(a) + \xi_{\circ 2}(a) + \dots + \xi_{\circ(j-1)}(a)$. This shows that for any $j \geqslant D$ it holds $\phi_{\circ j}(0) = \phi_{\circ d}(0)$. Since $\phi_{\circ M}(0) = 0$ for some $M \geqslant 1$, and thus we conclude that $\phi_{\circ j}(0) = 0$ for any $j \geqslant D$.

This implies that $\phi_{\circ j}(x) = \xi_{\circ j}(x)$ for any $j \geqslant D$ and any $x \in F$, and completes the proof of the lemma.

□

lemmaalongcoordinates

In Lemma 7.3 below we use additive notation for N and G/N.

lemmaalongcoordinates

Lemma 7.3. *Let G/N and N be Abelian groups. Let $w(x_1,\dots,x_n)$ be an identity of G/N and u and v be as in Remark 5.1, Assume that $G/N=\mathbb{Z}^d+A'$, A' is finite. Denote by $\epsilon_1,\dots,\epsilon_d$ the standard generators of $\mathbb{Z}^d$.*

i) There exists l_j ($j\in J$, J is a finite set), and $r_j=(r_{2,j},\dots,r_d)$ such that for any $x_2,\dots,x_n\in G$ and $x\in N$

$$\phi_u(x)=\sum_{j\in J, z=(r_{2,j}x_2,r_{3,j}x_3,\dots,r_{n,j}x_n)} l_j z(s)$$

and such that for any $k:1\leqslant k\leqslant d$ there exists $x_2,\dots,x_n\in G$ such that for any $x\in N$

$$\phi_u(x)=\sum_{j\in J} l_j x^{M_j\epsilon_k},$$

where M_j are integers such that $M_j\neq M_{j'}$ for $j,j'\in J$ such that $j\neq j'$.

In particular, if there exists $x_2,\dots,x_n$ such that the homomorphism ϕ_u from N to N is not identically zero, then there exists $x_2,\dots,x_n$ such that ϕ_u has the form as above and at least one of coefficients l_j is non-zero.

ii) Assume moreover that w is an iterated identity of G and that N contains at least one non-identity element. If there exists $x_2,\dots,x_n$ such that the homomorphism ϕ_u from N to N is not identically zero, then there exists $x_2,\dots,x_n$ such that ϕ_u has the form as above and at least two of coefficients l_j is non-zero.

Proof. We know that $w(x_1,\dots,x_n)=u(x_1,\dots,x_n)v(x_2,\dots,x_n)$, u and v are identities of G/N. Fix some x_2, …, x_n in G. Put $\psi(x)=\phi_u(x)=u(x,x_2,\dots,x_n)$. For any $x\in G$

$$u(x,x_2,\dots,x_n)=\prod\alpha_i(x_2,\dots,x_n)x^{l_i}\alpha_i(x_2,\dots,x_n)$$

Observe that for any $x\in N$ $\alpha_i(x_2,\dots,x_n)x^{l_i}\alpha_i(x_2,\dots,x_n)$ depends only on the image of α_i in G/N. Regrouping, if necessary, the terms with the same image of α_i in G/N, we conclude that for any $x_2,\dots,x_n\in G$ and $x\in X$ ϕ_u has the following form, in the additive notation on N and G/N:

$$\phi_u(x)=\sum_{j\in J, z=(r_{2,j}x_2,r_{3,j}x_3,\dots,r_{n,j}x_n)} l_j(s)^z$$

Now we fix $k:1\leqslant k\leqslant d$ and consider $x_2,\dots,x_n\in G/N$ (or to be more precise, consider $x_2,\dots,x_n$ in G with corresponding image in G/N $x_2=L_2\epsilon_k$, $x_3=L_2\epsilon_k$, $x_n=L_n\epsilon_k$. Consider the lexicographic order on $\mathbb{Z}^{n-1}$. Observe that if positive integers L_i are such that L_2 is much larger than L_3, L_3 is much larger than L_4, …($L_2>>L_3>>L_4>>\dots>>L_n$) and if $r=(r_2,\dots,r_n)>\bar r=(\bar r_2,\dots,\bar r_n)$ in the lexicographic order on $\mathbb{Z}^{n-1}$, then

$$L_2r_2+L_3r_3+\dots+L_nr_n>L_2\bar r_2+L_3\bar r_3+\dots+L_n\bar r_n$$

Thus, choosing a quickly decreasing sequence of positive integers $L_2,\dots,L_n$, we can assure that all $M_j=L_2r_{2,j}+L_3r_{3,j}+\dots+L_nr_{n,j}$, j belonging to the finite set J, are distinct.

This completes the proof of i).

Two prove ii) take some $x\neq e, x\in N$. Observe that if $\phi_u(x)=\sum_{j\in J} l_j x^{M_j\epsilon_k}$ for some choice of $x_2,\dots,x_n$ and all x and exactly one among coefficients l_j is non-zero, then any iteration of ϕ_u has the same property, which contradicts the fact that some iteration of $\phi_u(x)$ is trivial.

□

lemmacubeisfinite

Lemma 7.4. *Let H be an Abelian group, $A = \mathbb{Z}^d$ act on H and there exists $s \in H$ such that H is generated by $a(s)$, $a \in A$. Let $\epsilon_1, \dots, \epsilon_n$ denote the standard generators of $\mathbb{Z}^d$. Suppose that for any $i : 1 \leqslant i \leqslant d$ there exists $k > 0$ and integers $m_1^{(i)} < m_2^{(i)} < \dots m_k^{(i)}$ and $\gamma^{(i)}{}_1, \gamma_2^{(i)}, \dots, \gamma_{k_i}^{(i)}$ such that (for each i) at least two among $\gamma_j^{(i)}$ are non-zero and*

$$\sum_j \gamma_j^{(i)} s^{(m_j^{(i)}\epsilon_i)} = 0.$$

Then $F \otimes \mathbb{Q}$ has finite dimension.

Proof. Denote by π the tensor map from F to $F \otimes \mathbb{Q}$. For any $i : 1 \leqslant i \leqslant d$ chose $m_j^{(i)}$, $\gamma_j^{(i)}$ as in the assumption of the lemma. Without loss of generality we can assume that $\gamma_k^{(i)} \neq 0$, $\gamma_1^{(i)} \neq 0$ for each i. We have therefore

$$s^{m_{k_i}^{(i)}\epsilon_i} = -\frac{1}{\gamma_{k_i}^{(i)}} \sum_{j=1}^{k_i-1} s^{(m_j^{(i)}\epsilon_i)} \text{ and } s^{m_1^{(i)}\epsilon_i} = -\frac{1}{\gamma_1^{(i)}} \sum_{j=2}^{k_i} s^{(m_j^{(i)}\epsilon_i)}.$$

This implies that for any $a \in \mathbb{Z}^d$

$$s^{(m_{k_i}\epsilon_i+z)} = -\frac{1}{\gamma_{k_i}} \sum_{j=1}^{k_i-1} s^{(m_j^{(i)}\epsilon_i+z)} \text{ and } s^{(m_1^{(i)}\epsilon_i+z)} = -\frac{1}{\gamma_{k_i}} \sum_{j=2}^{k_i} s^{(m_j\epsilon_i+z)}.$$

Consider the finite cuboid $P \subset \mathbb{Z}^d$ containing $(z_1, \dots, z_d) : m_1^{(i)} \leqslant z_i \leqslant m_{k_i}^{(i)}$. Consider the finite set $p(s)$, $p \in P$, and denote this finite set by F_P. It is clear that the linear span of the image of F_P for the tensor mapping is equal to $F \otimes \mathbb{Q}$, and hence $F \otimes \mathbb{Q}$ is finite dimensional.

□

Let us say that a map $\phi : X \to X$ is weakly recurrent, if there exists a finite set $X_f \subset X$ such that for any x there exists $n \geqslant 1$ such that the n-th iteration $\phi_{\circ n}(x) \in X_f$.

Remark 7.5. If ϕi is recurrent (with respect to some fixed point $e \in X$), and if $d \geqslant 1$, then $\phi_{\circ d}$ is weakly recurrent.

Proof. Put $X_f = \{e, \phi(e), \phi_{\circ 2}(e), \dots, \phi_{\circ d}(e)\}$. It is clear, that if ϕ is recurrent, than for any x its iteration $\phi_{\circ d}$ visits the set X_f (infinitely many times). □

We return to the proof of the Theorem. We have some $x_2, \dots, x_n$ fixed, and we consider the dynamics of ϕ_u and ϕ_w.

Let T be the torsion subgroup of N. F/T is torsion free.

The homomorphism u preserves T and thus induces the quotient homomorphism $\phi_u^{(F/T)} : F/T \to F/T$. Let v' is the image of $v(x_2, \dots, x_n)$ in F/T. It is clear that

$$\pi_{F/T}(\phi_w(x)) = \phi_u^{(F/T)}(\pi_{F/T}(x)) + v'.$$

Since finitely generated metabelian groups satisfy maximal condition for normal subgroup (Hall, [19]), we know that N is a normal closure of a finite set S'. This implies that N is generated by s^a, $s \in S'$, $a \in A$ (where s^a denotes the action of a on s by conjugation). Considering the orbit of S under the action of the finite group A_f, we conclude that there exists a finite set $S \subset N$ such that N is generated by s^a, $s \in S$, $a \in \mathbb{Z}^d$

Let T be the torsion subgroup of N. Observe that T is a normal subgroup of G. Indeed, its conjugation by an element of G belongs to N since N is normal, and a conjugation of a torsion element is a torsion element.

hall1954

Therefore, using once again [19] we conclude that there exists a finite set S'_T such that T is generated as a normal subgroup by S'_T. In other words, T is generated by gsg^{-1}, where $g \in G$, $s \in S'_T$ and generated by asa^{-1}, where $a \in A$, $s \in S'_T$. Considering, as before, the orbit of S'_T under the action of the finite group A_f, we conclude that there exists a finite set $S_T \subset T$ such that T is generated by asa^{-1}, where $a \in \mathbb{Z}^d$, $s \in S_T$.

For any $s \in S$ consider the linear span V_s of the images $a(s), a \in \mathbb{Z}^d$ in $F \otimes \mathbb{Q}$. Let d_s denotes the dimension of V_s. Recall that S is a finite set. Let d be the maximum among finite values of d_s, $s \in S$. If all d_s are infinite, we put $d = 0$. Let us show that $\phi^{F/T}_{u,\circ d+1)}(x) = 0$ and $\phi^{F/T}_{w,\circ d+1}(x) = 0$ for any $x \in F/T$.

Observe that since ϕ_u commutes with the action of A and $\phi_u(s)$ belongs to the subgroup generated by $a(s)$, $a \in \mathbb{Z}^d$, it suffices to consider the action induces bt ϕ_u on each V_s. Observe that if $s \in S$ is such that the dimentsion d_s of V_s is $< \infty$, then for any $d > d_s$ $\phi^{F/T}_{u,\circ d}(t), \phi^{F/T}_{w,\circ d}(t) = 0$ for any $t \in V_s \cap \pi(F)$, and hence also for any $t \in V_s$ (follows from Lemma 7.2). Otherwise, if s is such that $d_s = \infty$, observe that ii) of Lemma 7.3 and Lemma 7.4 imply that the restriction of $\phi^{(F/T)}_u$ and $\phi^{(F/T)}_w$ to $V_s \cap \pi(F)$ (and hence the restrictions to V_s as well) are identitcally zero.

lemmafinitetensor lemmacubeisfinite lemmalongcoordinates

We have shown therefore that $\phi^{F/T}_{u,\circ d+1)}(x) = 0$ and $\phi^{F/T}_{w,\circ d+1}(x) = 0$ for any $x \in F/T$.

We implies that $\phi_{w,\circ d+1}(x) \in T$ for any $x \in F$. Now we consider $\bar{w} = w_{\circ d}$, chose $\bar{u}(x, x_2, \dots, x_n)$, $\bar{v}(x_2, \dots, x_n) \in T$ and consider the dynamics of the homomorphism $\phi_{\bar{u}}$ on T.

lemmadimfinitefinite

Lemma 7.6. *H is an Abelian group such that $m^k h = 0$ for all $h \in H$, where $m = p_1 \dots p_l$, p_i are prime. Suppose that $H \otimes \mathbb{Z}/p_i\mathbb{Z}$ (viewed as a vector space over $\mathbb{Z}/p_i\mathbb{Z}$) is finitely dimensional for all i. Then H is finite.*

Proof. First suppose that $l = 1$, H is an abelian group such that $p^k h = 0$ for all h. $H \otimes \mathbb{Z}/pZ$ is finitely dimensional. Consider $H_i = p^i H$. It is clear that H_i is a subgroup of H and $H = H_0 \supset H_1 \supset \dots \supset H_k = 0$. Observe that $H_0/H_1 = H/pH = H \otimes \mathbb{Z}/pZ$. By the assumption of the Lemma, this is a finitely dimensional vector space over a finite fields, therefore H_0/H_1 is a finite group. Observe that H_i/H_{i+1} is a quotient of H_0/H_1 by the map which is a multiplication by p^i, and hence H_i/H_{i+1} is finite for any i. This implies that H is finite.

Now consider the general case. Prove the statement by induction on l. The statement of the base of the induction $l = 1$ is already proven. Consider $H' = p_l^k H$. Observe that H/H' satisfies the assumption of the induction step for $l - 1$, and hence H/H' is finite. Observe also that H' satisfies the assumption of the statement of the induction base, and thus H' is finite. We conclude that H is finite.

□

We want to show that there exists a finite subgroup $T_f \subset T$ such that $\phi_{\bar{u},\circ k}(x) = 0$ for any $x \in T$. Here k is such that $m^k t = 0$ for any $x \in T$ and m is a product of (finitely many) distinct prime numbers.

Indeed, take $t \in T$ and consider T_t to be the linear span of t^a, $a \in A$ and T'_t to be the linear span of t^a $a \in \mathbb{Z}^d$. It is clear that T_t is equal to the linear span of $(T'_t)^a = T'_{t^a}$, $a \in A_f$. Observe T'_{t^a} is isomorphic to T'_t for any a, and hence the dimension of $T'_{t^a} \otimes \mathbb{Z}/p\mathbb{Z}$ is equal to that of $T'_t \otimes \mathbb{Z}/p\mathbb{Z}$ (and the same statement for the tensor product with $\mathbb{Q}$.)

If t is such that the dimension of $T_t \otimes \mathbb{Z}/pZ$ is infinite, then the dimension of $T'_t \otimes \mathbb{Z}/p\mathbb{Z}$ is infinite and also $T'_{t^a} \otimes \mathbb{Z}/p\mathbb{Z}$ is infinite for any $a \in A_f$. In this case Lemma 7.3 and Lemma 7.4 imply that all but possibly one coefficient of ϕ_u are divided by p. This implies

lemmacubeisfinite lemmalongcoordinates

that all such coefficients is divided by p, since otherwise there exist $x_2, x_3, \dots, x_n$ such that exactly one coefficients among l_j, defined in Lemma 7.3 is not divided by p. Observe that there exists $x \in T$ such that $x \notin pT$, and in this case any iteration of ϕ_u, applied to x, does not belong to pT, and this is in contradiction with the fact that one of such iterations is trivial.

Hence $\phi_{u,\circ k}(x)$, as well as $\phi_{w,\circ k}(x)$ belong to $m_t^k T$, where m_t is the product of p_i, such that p_i is a prime divisor of m and dimension of $T_t \otimes \mathbb{Z}/p_i\mathbb{Z}$ is infinite.

Now note that for any p_i the dimension of $m_t^k T \otimes \mathbb{Z}/p_i\mathbb{Z}$ is finite. Lemma 7.6 implies that the subgroup $m^k T$ is finite. Put T_f be the subgroup generated by $m_t^k T_t$, t is in the finite set S_T (the image of which under the action of A generates T). T_f is an Abelian group generated by a finite number of Abelian groups, and hence T_f is finite. Observe that $\phi_{u,\circ k}(t)$, as well as $\phi_{w,\circ k}(t)$ belongs to T_f for any $t \in T$.

We know therefore that there exists a finite group T_f and $C = D + k$ such that for any iterated identity w of G it holds $w_{\circ C}(x_1, \dots, x_n)$ belongs to T_f for any $x_1, \dots, x_n \in T_f$.

Now fix w and $x_2, \dots x_n \in G$. Let $B_w \subset T_f$ consists of the elements $z \in T_f$ such that $w_{\circ C'} e, x_2, \dots, x_n \in T_f$ for some $C : 0 < C'' \leqslant C$. Since w is an iterated identity, for any $x \in G$ there exists m such that $w_{\circ Cm}(x, x_2, \dots, x_n) \in B_w$. Any orbit of the word map $w_{\circ C}(*, x_2, x_3, \dots, x_n)$ has size at most $\#T_f$, and therefore for any $x \in G$ there exists $C_2 \leqslant \#T_f$ such that $w_{\circ CC_2}(x, x_2, \dots, x_n) \in B_w$. This implies that there exists $C_3 \leqslant C(\#T_f + 1)$ such that $w_\circ(x, x_2, \dots, x_n) = e$.

We have shown therefore that for any iterated identity w of G its iterational depth $s(w, G) \leqslant (D + k)(\#T_f + 1)$.

This completes the proof of the theorem.

□

8. S TYPE ITERATED IDENTITIES

Definition 8.1. We say that a group G satisfies S-type (or solvability type) iterated identity w if for any $x_1, \dots, x_n \in G$ there exists N such that for any $y_1, \dots y_{n^N}$ taking value among $x_1, \dots, x_n$ it holds

$$w_{\star N}(y_1, y_2, \dots, y_{n^N}) = w(w(w(\dots w(y_1, y_2, \dots y_n), w(y_{n+1}, y_{n+2}, \dots y_{2n}) \dots), \dots)) = e.$$

Example 8.2. $w_0(x_1, x_2) = x_1 x_2 x_1^{-1} x_2^{-1}$. *If G is a solvable group, then G satisfies S-type identity w_0.*

More generally, let $w(x_1, \dots, x_n)$ be such that the total number of each x_i in w is equal to zero. Then any solvable group G satisfies the S type identity w.

Proof. Let $G^{(n)}$ denotes the n-th group in the derived series of G, that is $G_1 = [G, G]$ and $G_{i+1} = [G_i, G_i]$ for any $i \geqslant 1$. Observe that w is an identity in any Abelian group, and hence $w(x_1, \dots, x_n) \in [G, G]$ for any $x_1, \dots, x_n \in G$. Moreover, if $x_1, \dots, x_n \in G^{(i)}$, then $w(x_1, \dots, x_n) in G^{(i+1)}$. Since G is solvable, there exists N such that $G^{(N)} = N$. Put $k = n^N$. It is clear that $w_{\star N}(y_1, \dots, y_k) = 0$ for any $y_1, \dots, y_i \in G$, □

Example 8.3. $\mathbb{Z}$ *satisfies an S type iterated identity $w(x_1, \dots, x_n)$ if and only if the total number of each x_i in w is equal to zero.*

Proof. Suppose that the total number of some x_i in w is equal to $m \neq 0$. Without loss of generality we can assume that $i = 1$. Let 1 denotes a generator of $\mathbb{Z}$. Consider a finite set consisting of 0 and 1. For each N put $k = 2^N$ and consider $y_1 = a$, $y_2 = y_3 = \dots y_k = 0$. Observe that $w_{\star N}(y_1, \dots, y_n) = m^N \neq 0$, and hence w is not an S type iterated identity of G.

Observe also, that if the total sum of each x_i in w is equal to zero, then w is an identity of $\mathbb{Z}$. This implies that w is an S type iterated identity of $\mathbb{Z}$. □

Example 8.4. *Let G be a solvable group containing at least one element of infinite order. Then G satisfies an S type iterated identity $w(x_1, \dots, x_n)$ if and only if the total number of each x_i in w is equal to zero.*

Proof. Follows from Examples 8.2 and 8.3. □

Similarly to E type iterated identities, the property to satisfy S type iterated identity is closed under taking subgroups, quotients, union of increasing group sequences, and it addition this property is closed under taking extensions:

Lemma 8.5. (1) *If G satisfies an S-type iterated identity w, then any subgroup of G has the same property.*
(2) *If G satisfies an S-type iterated identity w, then any quotient of G has the same property.*
(3) *If $G_1 \subset G_2 \subset G_3 \dots$, and each G_i satisfies an S-type identity w, then the union $G = \cup_{i=1}^{\infty} G_i$ satisfies the same property.*
(4) *If H is a normal subgroup of G, $H' = G/H$, H, H' satisfy an S type iterated identity w, then G has the same property*

Proof. (1)-(3) are straightforward.

To prove (4), observe that for any since G/H satisfies S-type iterated identity w, for any n_1 there exists N_1 such that for any $t_1, \dots, t_{n_1} \in G$ and any
$y_1, \dots, y_{k_1}$, $k_1 = n_1^{N_1}$ taking values among $t_1, \dots, x_{t_1}$

$$w_{\star N_1}(y_1, \dots, y_{k_1}) = w(w(w(\dots w(y_1, y_2, \dots y_{n_1}), w(y_{n_1+1}, y_{n_1+2}, \dots y_{2n_1}) \dots), \dots)) \in H.$$

Consider a set $V \subset H$ which consists of possible values of $w_{\star k} y_1, \dots, y_{k_1}$, where $k_1 = n_1^{N_1}$ and all y_i take value among $x_1, \dots, x_{n_1}$. It is clear that V is finite, since its cardinality, which we denote by n_2, is at most $n_1^{k_1}$.

Since H satisfies the S type iterated identity w, there exists N_2 (depending on V) such that for $k_2 = n_2^{N_2}$ and any $z_1, z_2, \dots, z_{k_2}$ taking values in V it holds

$$w_{\star N_2}(z_1, \dots, z_{k_2}) = w(w(w(\dots w(z_1, z_2, \dots z_{n_2}), w(z_{n_2+1}, z_{n_2+2}, \dots z_{2n_2}) \dots), \dots)) = e.$$

Put $N = N_1 + N_2$. Consider $k = n_1^N$. Observe, that for any $r_1, r_2, \dots, r_k$ taking values among $t_1, \dots, t_{n_1}$ it holds

$$w_{\star N}(r_1, \dots, r_k) = e,$$

and this concludes the proof of 4).

□

Given a class of groups C, denote by $E(C)$ the minimal class of groups, which is closed under taking four elementary operations, that is, taking subgroup, quotient, direct limit and group extensions.

Corollary 8.6. *Let w be a word Let $C_{w,N}$ be the variety of groups, generated by the identity $w_{\star N}$, that is, the class of groups that satisfy the identity $w_{\star N}$. Put $C_w = \cup_{N=1}^{\infty} C_{w,N}$. The class of groups satisfying an S type iterated identity w contains $E(C_w)$.*

In general, this class is larger than $E(C_w)$.

igorchuknotelementary

Example 8.7. *Let C be a class of groups of bounded 2-torsion. We know that the fist Grigorchuk group G_{Grig} satisfies Solvability type (as well as Engel type) iterated identity $w(x_1) = x_1^2$. However, G_{Grig} does not belong to $E(C_w)$.*

Proof. If the class C is closed with respect to taking subgroups and quotients, than it is not difficult to see that the elementary class $E(C)$ is the minimal class of groups, containing C and closed with respect to taking direct limits and extensions by a group in C (see Corollary 2.1 [23]).

josinetelementaryctasses

More precisely, given an ordinal α, denote by $E_\alpha(C)$ in the following way. Suppose we have defined $E_\beta(C)$ for any $\beta < \alpha$. If α is a limit ordinal, then $E_\alpha(C)$ is defined as a union $\cup_{\beta<\alpha}(C)$. If α is not a limit ordinal, then $E_\alpha(C)$ is the class of groups containing direct limits of groups in $E_{\alpha-1}$ and extentsions of groups in $E_{\alpha-1}$ by a group of C. Then for each $E_{\alpha-1}$, for any α, is closed with respect to taking subgroups and quotients (see Lemma 3.1 in [23]).

josinetelementaryctasses

Let $C = C_w$. It is clear that C is the class of groups of bounded 2-torsion and that C is closed with respect to takings subgroups and quotients.

Consider a minimal ordinal α such that E_α contains a G_{Grig}. It is clear that α is not a limit ordinal, and that G can be obtained by taking a quotient from the groups in $E_{\alpha-1}$ (since G is finitely generated, the last operation to obtain G could not be a direct limit). Therefore, there exists a normal subgroup N in G_{Grig} such that N and G/N belongs to $E_{\alpha-1}$. Since $G \notin E_{\alpha-1}$, $N \neq e$.

Any proper quotient of G_{Grig} is finite, G/N is finite, N is a normal subgroup of finite index in G. Any non trivial normal subgroup in G_{Grig} is of finite index and contains some congruence subgroup $St(n)$ (Grigorchuk, [?grigorchuk200][Proposition 10]). This implies that $St(n)$ belongs to $E_{\alpha-1}$. Since any congruence subgroup contains G_{Grig} as a quotient (taking restriction on the first branch of level n of the tree, [15]), this also implies that G_{Grig} belongs to $E_{\alpha-1}$. This is a contradiction that α is a minimal ordinal such that E_α contains G_{Grig}.

?grigorchuk200

?grigorchukparabolic

□

8.1. **Quasi-varieties.** Recall that a variety of groups is a class of groups for which a certain set of identities hold. This class is obviously closed under taking subgroups, homomorphic images (the same is to say: by taking quotients), and direct products.

Given a set Ω of iterated identities, we consider the class of groups G such that G satisfies an E type (correspondingly S type) iterated identity w, for all $w \in \Omega$. We call this class of groups an Engel type (correspondingly S type) *quasi-variety* generated by a set of iterated identities Ω.

Example 8.8. *Two nilpotent groups N_1 and N_2 generated the same E-type iterated quasi-variety if and only if they generated the same variety of groups and $m(N_1) = m(N_2)$, where $m(G) \in \mathbb{N} \cup +\infty$ (as defined in Example 5.9) is the minimal positive integer m such that such that for any element $g \in G$ there exists an integer $t > 0$ such that such that $g^{m^t} = e$ (and $+\infty$ if no such m exists).*

examplenilpotent

Example 8.9. *Any solvable group G containing at least one element of infinite order generated the same S type iterated quasi-variety as $\mathbb{Z}$*

We say that an E-type (correspondingly S-type) iterated identity w' is a corollary of an E-type (correspondingly S-type) iterated identity w if any group satisfying an E-type (correspondingly S-type) iterated identity w also satisfies w'. Similarly, given a set of iterated identities Ω and an iterated identity w we say that w is a corollary of Ω if any group satisfying all identities from Ω also satisfies w.

If we consider (usual) identities instead of iterated ones, this notion is not very interesting since it simply means that the subgroup corresponding to one of them contains the characteristic subgroup that corresponds to the other one. Given an iterated identity w, of E type or S type, it seems interesting to describe all corollaries of w. Similarly, given a set Ω of iterated identities, describe the corollaries of Ω.

openquestions

8.2. **Open questions.**

Question 8.10. Given a group G, describe the sets of iterated identities $\Omega_E(n)$ and $\Omega_S(n)$ that holds in this group (e.g. for Grigorchuk groups, for Golod groups, for Basilica group).

The same question one can ask for families of groups (finite or infinite). For example, what can be said about the sets of iterated identities $\Omega_E(n)$ and $\Omega_S(n)$ that hold for all solvable groups of the group G)? For examples of iterated identities w in the characterizations of finite solvable groups in Remark 2.6, what are finitely generated groups that satisfy the iterated identity w (which do not need to be solvable, see Remark 2.5)?

preformulationsolvable

remarkgolod

Question 8.11. For which groups G the characteristics subsets Ω_n of G are subgroups of F_n, for all n?

Question 8.12. Describe the E type and S type quasi-varieties generated by a group G (or by a family of groups).

Question 8.13. Which groups are quasi-free with respect to the iteration of identities?

Question 8.14. Are free Burnside group $B_{n,p}$on n generators bounded? Are free groups of a varieties of groups generated by a single identity bounded?

Question 8.15. For which sets $\Omega \subset F_k$ there exists a group G such that the set of iterated identities on k variables of the group G is equal to Ω?

Question 8.16. Let r be an odd prime number. For an integer n consider $w_n(x_1, x_2) = (x_1^{rn} x_2^{rn} x_1^{-rn} x_2^{-rn})^n$ Adyan shows that if we fix n large enough ($n \geqslant 4381$ in [2] and $n \geqslant 1003$ in [3][page 297]) then the following identities are independent , that is, none of them follows from the others.

adyan1970

adyanbook

Is it true that w_n , for n large enough, are independent both as Engel type iterated identities and Solvability type iterated identities?

REFERENCES

abert

[1] Miklós Abért, *Group laws and free subgroups in topological groups*, Bull. London Math. Soc. **37** (2005), no. 4, 525–534, DOI 10.1112/S0024609305004251X. MR2143732 (2006d:20005)

adyan1970

[2] S. I. Adjan, *Infinite irreducible systems of group identities*, Izv. Akad. Nauk SSSR Ser. Mat. **34** (1970), 715–734 (Russian). MR0286871 (44 #4078)

adyanbook

[3] S. I. Adyan, *Problema Bernsaida i tozhdestva v gruppakh*, Izdat. “Nauka”, Moscow, 1975 (Russian). MR0432770 (55 #5753)

aleshin

[4] S. V. Alešin, *Finite automata and the Burnside problem for periodic groups*, Mat. Zametki **11** (1972), 319–328 (Russian). MR0301107 (46 #265)

bandmanandco

[5] Tatiana Bandman, Gert-Martin Greuel, Fritz Grunewald, Boris Kunyavskiĭ, Gerhard Pfister, and Eugene Plotkin, *Identities for finite solvable groups and equations in finite simple groups*, Compos. Math. **142** (2006), no. 3, 734–764, DOI 10.1112/S0010437X0500179X. MR2231200 (2007d:20027)

manetalverbaldynamics

[6] Tatiana Bandman, Fritz Grunewald, and Boris Kunyavskiĭ, *Geometry and arithmetic of verbal dynamical systems on simple groups*, Groups Geom. Dyn. **4** (2010), no. 4, 607–655, DOI 10.4171/GGD/98. With an appendix by Nathan Jones. MR2727656 (2011k:14020)

ndmanetalsurjectivity

[7] Tatiana Bandman, Shelly Garion, and Fritz Grunewald, *On the surjectivity of Engel words on* $\mathrm{PSL}(2, q)$, Groups Geom. Dyn. **6** (2012), no. 3, 409–439, DOI 10.4171/GGD/162. MR2961281

fractalsets
[8] Laurent Bartholdi, Rostislav Grigorchuk, and Volodymyr Nekrashevych, *From fractal groups to fractal sets*, Fractals in Graz 2001, Trends Math., Birkhäuser, Basel, 2003, pp. 25–118. MR2091700 (2005h:20056)

brandlwilson
[9] Rolf Brandl and John S. Wilson, *Characterization of finite soluble groups by laws in a small number of variables*, J. Algebra **116** (1988), no. 2, 334–341, DOI 10.1016/0021-8693(88)90221-9. MR953155 (89g:20056)

braywilsonwilson
[10] John N. Bray, John S. Wilson, and Robert A. Wilson, *A characterization of finite soluble groups by laws in two variables*, Bull. London Math. Soc. **37** (2005), no. 2, 179–186, DOI 10.1112/S0024609304003959. MR2119017 (2005k:20035)

chou
[11] Ching Chou, *Elementary amenable groups*, Illinois J. Math. **24** (1980), no. 3, 396–407. MR573475 (81h:43004)

golod
[12] E. S. Golod, *Some problems of Burnside type*, Proc. Internat. Congr. Math. (Moscow, 1966), Izdat. "Mir", Moscow, 1968, pp. 284–289 (Russian). MR0238880 (39 #240)

grigorchukfirstgroup
[13] R. I. Grigorčuk, *On Burnside's problem on periodic groups*, Funktsional. Anal. i Prilozhen. **14** (1980), no. 1, 53–54 (Russian). MR565099 (81m:20045)

grigorchuk2000
[14] R. I. Grigorchuk, *Branch groups*, Mat. Zametki **67** (2000), no. 6, 852–858, DOI 10.1007/BF02675625 (Russian, with Russian summary); English transl., Math. Notes **67** (2000), no. 5-6, 718–723. MR1820639 (2001i:20057)

grigorchukparabolic
[15] Laurent Bartholdi and Rostislav I. Grigorchuk, *On parabolic subgroups and Hecke algebras of some fractal groups*, Serdica Math. J. **28** (2002), no. 1, 47–90. MR1899368 (2003c:20027)

gromov
[16] M. Gromov, *Hyperbolic groups*, Essays in group theory, Math. Sci. Res. Inst. Publ., vol. 8, Springer, New York, 1987, pp. 75–263. MR919829 (89e:20070)

guptasidki
[17] Narain Gupta and Saïd Sidki, *On the Burnside problem for periodic groups*, Math. Z. **182** (1983), no. 3, 385–388, DOI 10.1007/BF01179757. MR696534 (84g:20075)

guralnikplotkinshalev
[18] Robert Guralnick, Eugene Plotkin, and Aner Shalev, *Burnside-type problems related to solvability*, Internat. J. Algebra Comput. **17** (2007), no. 5-6, 1033–1048, DOI 10.1142/S0218196707003962. MR2355682 (2008j:20110)

hall1954
[19] P. Hall, *Finiteness conditions for soluble groups*, Proc. London Math. Soc. (3) **4** (1954), 419–436. MR0072873 (17,344c)

hallnilpotentgroups
[20] Philip Hall, *The Edmonton notes on nilpotent groups*, Queen Mary College Mathematics Notes, Mathematics Department, Queen Mary College, London, 1969. MR0283083 (44 #316)

magnus1935
[21] Wilhelm Magnus, *Beziehungen zwischen Gruppen und Idealen in einem speziellen Ring*, Math. Ann. **111** (1935), no. 1, 259–280, DOI 10.1007/BF01472217 (German). MR1512992

nekrashevychbook
[22] Volodymyr Nekrashevych, *Self-similar groups*, Mathematical Surveys and Monographs, vol. 117, American Mathematical Society, Providence, RI, 2005. MR2162164 (2006e:20047)

osinelementaryclasses
[23] D. V. Osin, *Elementary classes of groups*, Mat. Zametki **72** (2002), no. 1, 84–93, DOI 10.1023/A:1019869105364 (Russian, with Russian summary); English transl., Math. Notes **72** (2002), no. 1-2, 75–82. MR1942584 (2003h:20062)

pervova
[24] E. L. Pervova, *Everywhere dense subgroups of a group of tree automorphisms*, Tr. Mat. Inst. Steklova **231** (2000), no. Din. Sist., Avtom. i Beskon. Gruppy, 356–367 (Russian, with Russian summary); English transl., Proc. Steklov Inst. Math. **4 (231)** (2000), 339–350. MR1841763 (2002g:20051)

plotkinural
[25] B. I. Plotkin, *Notes on Engel groups and Engel elements in groups. Some generalizations*, Izv. Ural. Gos. Univ. Mat. Mekh. **7(36)** (2005), 153–166, 192–193 (English, with English and Russian summaries). MR2190949 (2006j:20053)

ribnere
[26] Evija Ribnere, *Sequences of words characterizing finite solvable groups*, Monatsh. Math. **157** (2009), no. 4, 387–401, DOI 10.1007/s00605-008-0034-6. MR2520689 (2011a:20034)

thompson1
[27] John G. Thompson, *Nonsolvable finite groups all of whose local subgroups are solvable*, Bull. Amer. Math. Soc. **74** (1968), 383–437. MR0230809 (37 #6367)

thompson456
[28] ———, *Nonsolvable finite groups all of whose local subgroups are solvable. IV, V, VI*, Pacific J. Math. **48** (1973), 511–592, ibid. 50 (1974), 215–297; ibid. 51(1974), 573–630. MR0369512 (51 #5745)

wilsonconditions
[29] J. S. Wilson, *Two-generator conditions for residually finite groups*, Bull. London Math. Soc. **23** (1991), no. 3, 239–248, DOI 10.1112/blms/23.3.239. MR1123332 (92k:20068)

zelmanov
[30] E. I. Zel′manov, *Solution of the restricted Burnside problem for 2-groups*, Mat. Sb. **182** (1991), no. 4, 568–592 (Russian); English transl., Math. USSR-Sb. **72** (1992), no. 2, 543–565. MR1119009 (93a:20063)

A.E.: C.N.R.S., DMA, ENS, PARIS, FRANCE